# CSA AEROSOL INSECTICIDE FORMULA SEVEN

#### FOR HOUSE AND GARDEN

#### **ACTIVE INGREDIENTS:**

\*Resmothrin [5-(phenylmethyl)
-3-furanyl]methyl 2,2-dimethyl-3(2-methyl-1-propenyl)cyclopropencearboxylate.....0.25%

INERT INGREDIENTS: ..... 99,75%

100.00%



\*Cis/trans isomers ratio max. 30%( $\pm$ ) cis and min. 70%( $\pm$ ) trans.

### KEEP OUT OF REACH OF CHILDREN

#### **CAUTION**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT


SHAKE WELL BEFORE USING

manufactured by CSA LIMITED, INC. HOUSTON, TX 77269-0347

NET WEIGHT: 12 OZ.

EPA REG. NO. 7056-180

EPA EST. NO. 7056-TX-1



Revised 2/23/93

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or sprzy mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. In the home, all food processing surfaces and utensile should be covered during treatment, or thoroughly weshed before we.

#### STATEMENT OF PRACTICAL TREATMENT

If in eyes or on skin: Flush with water. Seek medical attention if irritation persists.

#### **ENVIRONMENTAL HAZARDS**

This pesticide is highly toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marches, and potholes). Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

#### PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open fiame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54.5°C) may cause bursting. Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical e juipment because of possible shock hazards. Do not use on surfaces that can be harmed or stained by water.

## **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE:

Keep in a locked storage area. Keep away from heat and

open flame.

DISPOSAL: Replace cap and discard container in trash.

For use in and around hotels, motels, homes, greenhouses and lathranes; food handling establishments, packaging and processing mants, reed houses, commercial, industrial, office, institutional and storage buildings, saill, \* \* handling areas of milk, cheese and ice cream plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneliel, \* flour and feed mills, bakeries, bosts, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. \* Yentilate rooms for 2 hours before returning patients. Also for use in buses, automobiles, \* trucks, boxcars and ship cargo holes. Also for use in dog bannels and hoppe stables. In horse stables, not for application if homes will be used for \* \* food.

Shake well before each use. Remove protective cap. Aim spray opening away from person. Pask button to spray.

INDOORS

FOR HOUSE FLEE, MOSQUITORS, GNATS, FLYING MOTHS (Millers), FRUIT FLES A VINSGAR FLEE: Close all doors and windows. Point nozzie upward; direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

FOR CRAWLING INSECTS: COCKROACHES (Waterbays), EARWIOS, GRAIN MI I'ES, SOWBUGS (Pill bage), SILVERFISH (Firebrate), SPIDERS, CENTIPEDES, MILLIPEDES, PLEAS, ANTS, BOOKLICE AND CRICKETS: Spray thoroughly and frequently into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, door frames, bookcases and storage areas. Spray directly on insects where possible. Repeat as accessary.

TO KILL ANTS - INCLUDING FIRE ANTS: Spray doors, around window frames, and other points of cutry. Spray ant trails, nosts, and hills contacting as many ants directly as possible.

TO EILL BEDBUGS AND LICE: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bed boards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may hard-or pests, concentrating on tuffs, folds, and seems. Repeat as necessary. For persistent infestations consult professional applicator.

with KILL FLEAS AND TICKS ON DOGS, CATS AND PREMISES: Apply directly nimals along back, under forelegs, and on other areas except scrotum and anus. Part hair to contact skin and insure good coverage. Hold nozzle about 1 foot (30 cm) away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes or mouth of animal. Repeat within 24 hours of first treatment, if necessary. To reduce reinfestation, thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30 to 46 cm). Also spray the entire inside surface of dog houses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible.

TO PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, begonias, geraniums and coleus for control of aphids, exposed thrips, adult whiteflies, spider mites, and fungus grats. Use with caution. To ascertain possible phytotoxicity, spray only a few plants, wait 24 boars, and check for possible injury.

FOR PROTECTION OF WOOLENS: Apply only to garments, blankets, carpets, and other woolen fabrics before placing in storage for protection from larvae of webbing clothing moths and black carpet beetles. Dry clean or wash garments and blankets before treating. Clean carpets, draperies and other fabrics by thoroughly brushing before treating. Hold the container 14 to 18 inches (30 to 46 cm) from the fabric and apply the spray while moving evenly in a forward and back motion across the fabric while covering one foot (30 cm) per second. Spray the entire fabric surface watil it is slightly moist or damp to the touch. When possible treat both sides of the fabric. Do not sock or wet fabric. For maximum protection place treated fabrics after drying in plastic or paper bags. Also treat cracks and crevices of closets, chosts and trunks where these fabrics are being stored. Treat the cracks and crevious at the rate of 2 seconds per linear foot (30 cm). When applied as directed on fabric and in storage areas, this treatment will offer protection for 6 months. To insure protection, inspect the fabric in the storage area ours, every three mouths. Repeat application only when necessary. Avoid heavy applications. When possible, make applications outdoors. L. application is made indoors, do not remain in treated areas. Ventilate the area afte: treatment is complete.

#### **OUTDOORS**

TO KILL FLYING INSECTS: Spray areas infested with mosquitoes, gasts (midges), house flies, stable flies, deer flies and flying moths (millers). Follow precautions listed above. Hold container 18 to 24 inches (46 to 61 cm) from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per

cubic yard (0.76 cm. m) of space while moving away from the trusted areas. To reduce annoyance from these flying insects, treat open porches and paties in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

TO PROTECT ORNAMENTAL. ANTS: To contral aphids, exposed theirs, adult white flies, leafhoppers, gypsy mot. s, plant bugs, spittlebugs and Japanese besties on smaller rhododendrous, pines, spruce, homlock, spires, dogwood, paics, mimoss, toyan, viburnum, succeymus, cales, meples, locust, forsythis and althou, and mimoss webworm leaves on missons, spray directly on insects. Hold container 18 inches (46 cm) from the infected plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot of growth per second Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control. Spray poonies, iris, zinnia snapdragous, ageratums, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphno. fatshedera, ivy, tobira, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above.

Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cotoneaster Rose, Camellia, Azalea, Pyracantha and Caenothus may also be treated to control the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

TO KILL TENT CATERPILLARS: Application should be made when caterpillars are young and tents are first noticed (usually late spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

TO KILL GYPSY MOTH CATERFILLARS: Application should be made when caterpillars are young and first noticed (usually late spring). Spray plants evenly to the point of run-off applying to both sides of leaves and other areas of plants where insects are suspected. For dense evergreens and shrubs, be sure spray reaches interior leaves and branches.

TO CONTROL NUISANCE PESTS: CRICKPTS, SELVERFISH, EARWIGS, SOWBUGS, CENTEPEDES AND MELLIPROES: Ser ; as directed above for indoor use.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FIRE ANTS: Kills and temporarily reduces annoyance of these interest. To reduce annoyance from these auts and keep them from entering houses, spray pation, barbecue areas, driveways, door sills, window sills, other elargice, areas, foundations of buildings, and trails, around pools and areas where the ants are noticed. Spray until the area is wet contacting us grany insects as possible. This treatment will not kill the colonies, but will reduce the immediate population for up to four hours. Repeat a pecus area. Avoid contacting plants, shrubbery and large areas of iswas.

FLEAS AND TICKS: Same as directed for indoor use, \* • \*

SPIDERS: (Including black Widow and Brown Recluse): Direct short bursts of spray on spiders and repeat as necessary.

WASPS, HORNET, YELLOW JACKETS AND BEES: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

