Sterilizing and Disinfecting Solution

Rapid-acting / Non-staining / Non-rusting

Sporicidal

Tuberculocidal

Virucidal :

Bactericidal

Fungicidal

Pseudomonacidal



long-life activated dialdehyde

CC-850

Artorook

Sterilizing and Disinfecting Solution

Rapid-acting Non-staining Non-rusting

Sporicidal

Virucidal

Fungicidal

Tuberculocidal

Bactericidal

Pseudomonacidal

CCCCONCENTRATE *

long-life activated dialdehyde

ong-life activated dialder.yde

Activator Vial
For Use With CIDEX Solution

Product CC-850

Active Ingredient
Glutaraldehyde ... 6.0%
Inert Ingredients ... 94.0%

Total .. 100.0%

Contains Sodium Nitrite as a corrosion inhibitor.

Note — Attached vial contents *must* be added to solution before this product is effective. See directions.

Rinse empty container thoroughly with water prior to discarding.

SEP 26 1975

CONCIDENTIAL ACCENTICIDE OF THE POLICY OF TH

DANGER: See Back Label for Precautions

*TRADEMARK

t See Directions for Use

Arbrook*

Directions for Dilution and Activation:

Add contents of this container of CIDEX 3X Concentrate to 2 gallons of distilled or deionized water. Then slowly add contents of attached vial. Stir well. Solution is ready to use when salts are completely dissolved.

Directions for Use:


For complete sterilization or disinfection of medical instruments and appliances, thoroughly clean, rinse and rough-dry objects before immersing in CIDEX Solution. Cleanse and rinse the lumens of hollow instruments before filling with CIDEX Solution.

Disinfection and Terminal Decontamination: Immerse completely for a minimum of 10 minutes to destroy vegetative pathogens including Mycobacterium tuberculosis, Pseudomonas aeruginosa and viruses (Poliovirus Type 1, 2, 3; Influenza Type A₁, A₂ (Hong Kong), B; Vaccinia; Coxsackievirus B-1; Reovirus Type 3; Herpes simplex) on inanimate surfaces

Sterilization: Immerse completely for a minimum of 10 hours to destroy resistant pathogenic spores including Clostridium sporogenes and Clostridium tetani. Remove instruments from CIDEX Solution using sterile technique and rinse thoroughly with sterile water

When activated, solution changes color to green.

Do not use activated stock solution beyond 28 days after mixing.



DANGER:

FOLLOW DIRECTIONS FOR USE

HARMFUL IF SWALLOWED

Skin — Avoid skin contact, as possibility of sensitization exists
May cause skin irritation. For skin contact flush thoroughly with water
Eyes — Contact with eyes causes damage. In case of contact
flush with water immediately and get medical attention

Food — Avoid contamination of food

Inhalation — Use in ventilated area

Instruments — CIDEX Solution will not prevent or remove oxidation caused by poor or improper plating, cleaning, or care of instruments.

KEEP IN A COOL PLACE

Especially recommended for terminal decontamination procedures as well as sterilization or disinfection of . . .

Lensed instruments
Anesthesia equipment
Respiratory therapy equipment
Rubber objects

Plastic objects
Sharp instruments
Thermometers

1 Dido . N

#il.

CONTENT ONE GAL (128 FLUID OZ)
MADE IN USA.
PAT. 3016328
CONTROL NUMBER ON BOTTOM

Arbrook inc.
Arlington, Jexas 76010

