#### CONCENTRATE PRODUCT

IN GAL PUT-UP

### CENTER PANEL

Sterilizing and Disinfecting Solution

Rapid-acting/nonstaining/nonrusting
Virucidal Tuberculocidal
Fungicidal Bactericidal
Sporicidal Pseudomonicidal

ESPECIALLY RECOMMENDED FOR REPEATED USE IN BUCKET AND TRAY SYSTEMS.

CIDEX\* II concentrate

Activated Dialdehyde

Active Ingredient
Glutaraldehyde--- 12.0%
Inert Ingredients---- 88.0%
TOTAL 100.0%

Contains sodium nitrite as a corrosion inhibitor.

NOTE: Activator must be added to diluted fluid before this product is effective. See directions.

KEEP OUT OF REACH OF CHILDREN

DANGER: See right Panel for precautionary statements.

\*See Directions for use.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

MAR 2 2 1983

\*Trademark of SURGIKOS, Inc.

Under the provided for mode, Provided to the first the posterior restricted under EPA Reg. So

RECEIVED

FEB 2 , 1983

Registration Division (TS-Z6Z)

#### CONCENTRATE PRODUCT

#### IN GAL PUT-UP

## LEFT PANEL

GENERAL CLASSIFICATION- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Reorder No.

#### DIRECTIONS FOR ACTIVATION:

- Prior to activation, dilute contents of one-gallon container of CIDEX\* II Fluid Concentrate with four (4) gallons of water.
- 2. After diluting, add contents of one 22-ounce container of Activator to solution.
- 3. Mix; solution is ready to use. When activated and diluted CIDEX II Solution changes color to yellow.

#### DIRECTIONS FOR USE:

For complete sterilization or disinfection of medical instruments and equipment, thoroughly clean, rinse and rough-dry objects before immersing in CIDEX II Solution. Cleanse and rinse lumens of hollow instruments before filling with CIDEX II Solution.

DISINFECTION AND TERMINAL DECONTAMINATION: Immerse completely for a minimum of 10 minutes to destroy vegetative pathogens including Mycobacterium tuberculosis, Pseudomonas aeurginosa and Viruses [Poliovirus Type I; Influenza Type A2 (Hong Kong); United to the control of the c

STERILIZATION: Immerse completely for a minimum of 10 hours to destroy resistant pathogenic spores including Clostridium sporogenes and Bacillus subtilis. Remove instruments from CIDEX II Solution using sterile technique and rinse thoroughly with sterile water.

Solution may be repeatedly used for not more than 14 days.

EPA Reg. No. \_\_\_\_\_EPA Est. No. \_\_\_\_

ACCEPTED
with COMMENTS
in EPA I, ther Dated:

Form de and Rodent ente Act us anacided, for the positione registered upder EPA Reg. No.

\*Trademark SURGIKOS 1982

RECLA

FES 2 3 1323

Registration Decade (15/10/)

## RIGHT PANEL

Especially recommended for disinfection of...

Anesthesia equipment Rubber objects
Respiratory therapy equipment Plastic objects
Metallic equipment or instruments Glass objects

STORAGE AND DISPOSAL

Keep in a cool place. Triple rinse empty container with water and dispose in an incinerator or landfill approved for pesticide containers or bury in a safe place.

Content 128 fl. oz. (3.785 liters) Made in USA Control number on bottom

ACCEPTED
with COMMENTS
In EPA Letter 10 and

į

PRECAUTIONARY STATEMENTS

MAR 2 2 1333

DANGER:

HAZARD TO HUMANS FOLLOW ALL LABEL DIRECTIONS Under the state of the state. Further the Art is a state of the transportation for the perturbed register to a september 1,000 Per 120 Per 120

Prevent contact with skin or eyes by using eye protection and rubber gloves when handling or pouring.

STATEMENT OF PRACTICAL TREATMENT

EYES-- Corrosive, causes eye damage. Do not get in eyes. In case of contact, flush with water immediately and get medical attention.

SKIN-- Avoid skin contact, as possibility of skin sensitization exists. Can cause skin irritation. For skin contact, flush thoroughly with water. Get medical attention if irritation occurs.

INGESTION-- Harmful if swallowed. Avoid contamination of food.

INHALATION-- Avoid spillage, use in a well-ventilated area.

RECEIVING

FEB 2 3 1983

Registration Division (TS-767)

## ACTIVATOR LABEL



BANGER Consert with eyes causes dimage in case of sensect with eyes fault only and get medical enterior. For sun sensect, both the terror and an extract.


Area empty container thereughly

# **ACTIVATOR**

For use with CIDEX\* II Fluid only

TRANSPORTE VED





RECEIVA

. 3

FEB 2 3 1983

Registration Division ((S-767)

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (NH-567) MASSINGTON LIC 1, 4/	TERM OF ISSUANCE	22 MAR 1983
NOTICE OF PESTICIDE: 1 THE SETTING THE	NAME OF PESTICIDE PRODUCT	
(Under the Tederal Insecticide Tuma (ph) and Rodentica be Act (a) are indeal.	7	
AME AND ADDRESS OF REGISTRANT (Include 71P code)	<del></del>	
Г	┑,	
1	ı	
<b>L</b>	٦	
NOTE: Changes in labeling formula differing in substance fr submitted to and accepted by the Registration Division prior product always refer to the above U.S. EPA registration num	r to use of the label in commerce, ber,	In any correspondence on
On the basis of information furnished by the registrant, the a the Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is hereby t	Registered/Reregistered u
A copy of the labeling accepted in connection with this Reg	gistration/Reregistration is return	ed herewith.
Act is not to be construed as giving the registrant a right to by others.		
•		
ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		

1\_ .=

÷