

Front Panel

SURFACIDE/6
DETERGENT/DISINFECTANT

ACTIVE INGREDIENTS
 n-Alhyl (60% C₁₂, 30% C₁₄, 5% C₁₆, 5% C₁₈) dimethyl benzyl ammonium
 chlorides 4.50%
 n-Alhyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium
 chlorides 4.50%

INERT INGREDIENTS 91.00%

TOTAL 100.00%


EPA REG. NO. 7301-10
 EPA EST. NO. 7301 IL 01

DANGER:
KEEP OUT OF REACH OF CHILDREN.
 See outer box for practical treatment and additional
 precautionary statements, storage and disposal
 information, plus complete use directions.

manufactured by
 Highland Park, Ill. 60035

FOR INSTITUTIONAL/INDUSTRIAL USE ONLY.
ONLY FOR SALE TO, USE, AND
STORAGE BY SERVICE PERSONS.

NET WEIGHT 1000 g



Back Panel

DIRECTIONS FOR USE

DISINFECTION: To disinfect hard inanimate surfaces (such as walls, floors, table tops) use 1 ounce (1 packet) per gallon of water. Apply solution with mop, cloth, sponge or mechanical sprayer. Wet thoroughly. Allow surface to remain wet for 10 minutes, then let air dry.

To disinfect toilet bowls — flush toilet, add 1 ounce (1 packet) directly to bowl water. Swab bowl completely using scrub brush or toilet mop. Make sure to scrub under rim. Let stand for 10 minutes, then flush.

For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

MILDEWSTAIN: To control mold and mildew on pre-cleaned hard inanimate surfaces use 1 ounce (1 packet) per gallon of water. Apply solution with cloth, sponge, mop or mechanical sprayer. Wet thoroughly. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears.

CONTAINER DISPOSAL: Do not reuse empty packet. Wrap empty packet and put in trash collection.

SURFACIDE/6®

DETERGENT/DISINFECTANT

Deodorizer Saptizer Virucide Fungicide (against
pathogenic fungi) Mildewstat (on hard inanimate surfaces)

ACTIVE INGREDIENTS:

n-Alkyl (80% C₁₂, 30% C₁₄, 5% C₁₆, 5% C₁₈)

dimethyl benzyl ammonium chlorides

n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl

ethylbenzyl ammonium chlorides

INERT INGREDIENTS

4.5%
4.5%
91.00%
Total 100.00%

DANGER:

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautionary
and practical treatment statements. Keep
packets in box until ready for use.

EPA REG. NO. 7101-10

EPA EST. NO. 7101-IL-01

CONTENTS: 25 1-oz. packets
(29.6 ml each)

NET WEIGHT: 25 fl. oz.
(1 lb. 9 oz./739 ml.)




manufactured by
walton-march inc.

HIGHLAND PARK, ILLINOIS 60030

FOR INSTITUTIONAL/INDUSTRIAL USE ONLY
ONLY FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS.

Hazard to Humans and Domestic Animals



DIRECTIONS FOR USE
General Classification

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SURFACIDE/6 is designed for hospitals, nursing homes, schools, food processing plants and other institutions where housekeeping is of prime importance in controlling the hazard of cross-contamination.

DISINFECTION: To disinfect hard inanimate surfaces (such as walls, floors, table tops), use 1 ounce (1 packet) per gallon of water. Apply solution with mop, cloth, sponge or mechanical sprayer. Wet thoroughly. Allow surface to remain wet for 10 minutes, then let air dry.

To disinfect toilet bowls — flush toilet, add 1 ounce (1 packet) directly to bowl water. Swab bowl completely, using scrub brush or toilet mop. Make sure to scrub under rim. Let stand for 10 minutes, then flush.

For heavily soiled areas, a pre-cleaning step is required. Prepare fresh solution for each use.

MILDEWS/6: To control mold and mildew on pre-cleaned, hard non-porous surfaces, use 1 ounce (1 packet) per gallon of water. Apply solution with cloth, sponge, mop or mechanical sprayer. Wet thoroughly. Let air dry. Prepare fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

SANITIZATION: To sanitize previously cleaned and rinsed equipment or non-porous food contact surfaces, use 1 ounce (1 packet) per 3.5 gallons of water. **NOT POTABLE** Rinse thoroughly at this level. Apply with cloth, sponge, mop or mechanical sprayer. Wet thoroughly. Let stand for 10 minutes. Let air dry. Prepare fresh solution for each use.

7/8/70