

IMPORTANT

Baits should be examined regularly and replenished before completely eaten. Cereal baits prepared with Pival are resistant to insect infestation, less susceptible to mold and highly acceptable to rats and mice. However, should baits become sour, moldy or infested with insects, they must be replaced.

Do not reuse container. Destroy when empty.

CAUTION

NOTE: FOR PHYSICIANS: Intravenous injections of Vitamin K1 is indicated together with administration of blood transfusions.

EPA Reg. No. 7122-8

Date

Batch No.
AR-212

Packed By

ACCEPTED
11/13

UNDER THE FEDERAL FOOD, DRUG AND COSMETIC ACT
FOR ECONOMY AND EFFICIENCY
NEED NO

11/22/58
W. H. ...

GUARDIAN



Net
Contents:

PIVAL CONCENTRATE C. F.
A RODENTICIDE

ACTIVE INGREDIENT		
2 Pivalyl-1,3-indandione*		0.5%
INERT INGREDIENTS		99.5%
	TOTAL	100.0%

(See side panel for additional cautions.)

Or Chem Corp

1514 Eleventh Street
Portsmouth, Ohio 45662
Phone 614/353-1125

DIRECTIONS

Mix thoroughly one part of PIVAL with nineteen parts of a suitable bait material, such as corn meal, rolled oats or other cereal grains. A small amount of an edible oil such as mineral, corn or peanut oil may be used up to 5% to prevent dusting or to increase acceptability.

GENERAL DIRECTIONS

- a) Place baits in locations where rats and mice feed, water, and travel.
- b) Bait placement along walls, in corners and in concealed locations where rats and mice can feed unobserved are desirable.
- c) Contaminated or spoiled baits should be replaced promptly.

DIRECTIONS FOR RAT CONTROL

- a) Use 4 to 16 ozs of bait per placement.
- b) An uninterrupted supply of bait should be maintained for at least 10 days.

DIRECTIONS FOR MOUSE CONTROL

- a) Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed 3 to 12 foot intervals.
- b) An uninterrupted supply of bait should be maintained for at least 15 days.

Baits must be protected from dogs, cats, poultry, etc. by bait boxes, cage or other suitable means. For complete control, baiting should be maintained continuously as long as any evidence of feeding is observed. Average kill times are from 6 to 14 days, but some feedings may continue up to the fourth week. Where reinfestation is likely, permanent bait stations should be used and bait placed in them as often as necessary.

IMPORTANT

Baits should be examined regularly and replenished before completely eaten. Cereal baits prepared with Pival are resistant to insect infestation, less susceptible to mold and highly acceptable to rats and mice. However, if baits become sour, moldy or infested with insects they must be replaced.

Do not reuse container. Destroy when empty.

NOTE: FOR PHYSICIANS: Intravenous injections of Vitamin K1 is indicated together with administration of blood transfusions.

EPA Reg. No. 7122-8

Date

Batch No.

AR 212

Packed By

FACTORY

GUARDIAN



Net
Contents

PIVAL CONCENTRATE C. F.

A RODENTICIDE

ACTIVE INGREDIENT

2 Pivalyl 1,3 ind ind. 75

INERT INGREDIENTS

33.5

TOTAL

108.5

See side panel for additional cautions.

Ar Chem Corp

1514 Eleventh Street
Portsmouth, Ohio 4662
Phone 614/353-1125

DIRECTIONS

For general directions, see side panel. For specific directions, see side panel.

GENERAL DIRECTIONS

For general directions, see side panel. For specific directions, see side panel.

DIRECTIONS FOR RAT CONTROL

For directions for rat control, see side panel.

DIRECTIONS FOR MOUSE CONTROL

For directions for mouse control, see side panel.