BEEF CATTLE: To control lice, ticks and hornflies, use at the rate of 2 quarts per 50 gallon of water. Apply as a complete coverage spray, Repeat every two weeks if necessary. DO NOT APPLY TO LACTATING DAIRY ANIMALS.

POULTRY: To control Northern Fowl Mite, Poultry Lice and as a supplement to premise treatments for Chicken Red Mite use 2 tablespoons per 1 gallon of water. Spray 1 gallon of dilute mixture per 100 to 150 binds.

POULTRY HOUSES: To control Northern Fowl Mite, Chicken Red Mite, Poultry Lice, Flies and Mosquitoes, use 4 tablespoons per 1 gallon of water. Apply to walls, ceilings, roosts, nests, and cracks and crevices. Repeat as necessary.

FLIES: Use at the rate of 2 quarts per 25 gallon of water (1 pint to 6 gallon). For best results add one pound of sugar for each gallon of finished spray. Mix well, Spray all surfaces and places where flies and mosquitoes light or congregate. Use a wetting spray and spray to the point of run-off. Use around buildings, stockyards, dairy barns, poultry houses, garbage cans, and manure piles. Avoid contamination of feed and foodstuffs, Repeat as necessary.

FLOWERS, SHRUBS AND SHADE TREES: Use at the rate of 1 quart per 100 gallons of water (2 teaspoons per gallon). Mix well and spray plants thoroughly. This spray will kill aphids, red spider mites, bagworms, leaf-hopper, thrips, scales, meallybugs, white flies, leaf miners, leaf bugs, adult Japanese beetles, lace bugs, plantbugs and tent caterpillars. Repeat treatment as necessary. Make applications for scales when crawlers settle. CAUTION: INJURY MAY OCCUR TO CER-TAIN TENDER PLANTS SUCH AS CERTAIN FERNS AND SOME SPECIES OF CRASSULA. DO NOT TREAT THESE.



Malathion WE

Emulsifiable Concentrate 50%

Product No. 6202

Active Ingredients: Malathion O.O-Dimethyl dithio-phosphate of diethyl mercapiosuccinate)
Aromatic petroleum hydrocarbons
Inert Ingredients: TOTAL

CAUTION

Keep Out Of Reach Of Children See side panel for other cautions;



1514 Eleventh Street Portsmouth, Ohio 45662 Phone 614/353-1125

HOUSEHOLD INSECTS: For control of Roaches, Ants. Spiders. Silverfish, Crickets, Clover Mites and Earwigs, use at the rate of 3 tablespoonfuls per quart of water (1 quart per 5 gallons). Apply as a coarse spray or with a paint brush to window sills, baseboards, drainboards, under sinks, and stoves and other areas frequented by insects.

AVOID CONTAMINATION OF FEED, FOOD UTENSILS, MILK, MILKING EQUIPMENT AND WATER.

DO NOT USE IN EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERCIALLY PREPARED OR PRO-CESSED, DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSED.

CAUTION

Harmful if swallowed. Avoid contact with skin; wash thoroughly with warm water and soap after using. Avoid contamination of feed or food stuffs, KEEP OUT OF REACH OF CHILDREN.

External: In case of spillage on skin, wash with warm water

Internal: Atropine is antidote. CALL A PHYSICIAN IM-MEDIATELY.

This product is POISONOUS to fish, Keep Out of Any Body of Water.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

DO NOT USE ON PETS OR WILDLIFE.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Container Disposal

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

EPA REG. NO. 7122-42

DATE

ACCEPTED

7122-42