

DIRECTIONS FOR USE

BEE, CATTLE: For control of ticks and chiggers use at the rate of 1/2 quart per 100 gallons of water. Apply to cattle and sheep every 10 days. Repeat as necessary. DO NOT APPLY TO LACTATING DAIRY ANIMALS.

POULTRY: For control of Northern Fowl Mite, Poultry Lice and Red Mite. Poultry Lice, Fowl Mites and Mosquitoes. Use at the rate of 2 quarts per 25 gallons of water. Apply to all surfaces, including roosts, nests, and cracks and crevices. Repeat as necessary.

POULTRY HOUSES: For control of Northern Fowl Mite, Red Mite, Poultry Lice, Fowl Mites and Mosquitoes. Use at the rate of 2 quarts per 25 gallons of water. Apply to all surfaces, including roosts, nests, and cracks and crevices. Repeat as necessary.

FLIES: Use at the rate of 2 quarts per 25 gallons of water plus 1/2 gallon of kerosene. For best results add one pound of sugar for each gallon of finished spray. Mix well. Spray all surfaces and places where flies and mosquitoes light or congregate. Use a wetting spray and spray to the point of run-off. Use around buildings, stock pens, barns, poultry houses, garbage cans, and manure piles. Avoid contamination of feed and foodstuffs. Repeat as necessary.

FLOWERS, SHRUBS AND SHADE TREES: Use at the rate of 1 quart per 100 gallons of water. 2 teaspoons per gallon. Mix well and spray plants thoroughly. This spray will kill aphids, red spider mites, bagworms, leaf-hopper thrips, scales, mealybugs, white flies, leaf miners, leaf bugs, adult Japanese beetles, lace bugs, plantbugs and tent caterpillars. Repeat treatment as necessary. Make applications for scales when crawlers settle. **CAUTION: INJURY MAY OCCUR TO CERTAIN TENDER PLANTS SUCH AS CERTAIN FERNS AND SOME SPECIES OF CRASSULA. DO NOT TREAT THESE**

GUARDIAN



Premium Grade

Malathion WE Emulsifiable Concentrate 50%

Product No 6202

Active Ingredients

Inert Ingredients

TOTAL

Or Chem Corp

1514 Eleventh Street
Portsmouth, Ohio 45662
Phone 614/353-1125

EPA Reg. No. 7122-42
DATE 6-27-72

Batch

Packed by

Add to Food and Drug Administration

Use at the rate of 2 quarts per 25 gallons of water

plus 1/2 gallon of kerosene. For best results add one pound of sugar

for each gallon of finished spray. Mix well. Spray all surfaces and places

where flies and mosquitoes light or congregate. Use a wetting spray and spray

to the point of run-off. Use around buildings, stock pens, barns, poultry houses,

garbage cans, and manure piles. Avoid contamination of feed and foodstuffs.

Repeat as necessary.

Use at the rate of 1 quart per 100 gallons of water. 2 teaspoons per gallon.

Mix well and spray plants thoroughly. This spray will kill aphids,

red spider mites, bagworms, leaf-hopper thrips, scales, mealybugs,

white flies, leaf miners, leaf bugs, adult Japanese beetles, lace bugs,

plantbugs and tent caterpillars. Repeat treatment as necessary.

Make applications for scales when crawlers settle.

CAUTION: INJURY MAY OCCUR TO CERTAIN TENDER PLANTS SUCH AS CERTAIN FERNS AND SOME SPECIES OF CRASSULA. DO NOT TREAT THESE

DIRECTIONS FOR USE

BEEF CATTLE

For control of ticks use at the rate of 2 quarts per 25 gallons of water (1 quart per 6 gallons). For best results add one pound of sugar for each gallon of finished spray. Mix well. Spray all surfaces and places where ticks light or congregate. Use a wetting spray and spray to the point of run-off. Use around buildings, stockyards, dairy barns, poultry houses, garbage cans, and manure piles. Avoid contamination of feed and foodstuffs. Repeat as necessary.

POULTRY

For control of Northern Fowl Mite, Poultry Lice and Red Mite use at the rate of 2 quarts per 25 gallons of water. Spray 1 gallon of finished spray per 25 gallons of water.

POULTRY HOUSES: To control Northern Fowl Mite, Chicken Red Mite, Poultry Lice, Flies and Mosquitoes, use 4 tablespoons per 1 gallon of water. Apply to walls, ceilings, roosts, nests and cracks and crevices. Repeat as necessary.

FLIES: Use at the rate of 2 quarts per 25 gallons of water (1 quart per 6 gallons). For best results add one pound of sugar for each gallon of finished spray. Mix well. Spray all surfaces and places where flies and mosquitoes light or congregate. Use a wetting spray and spray to the point of run-off. Use around buildings, stockyards, dairy barns, poultry houses, garbage cans, and manure piles. Avoid contamination of feed and foodstuffs. Repeat as necessary.

FLOWERS, SHRUBS AND SHADE TREES: Use at the rate of 1 quart per 100 gallons of water (2 teaspoons per gallon). Mix well and spray plants thoroughly. This spray will kill aphids, red spider mites, bagworms, leaf-hopper, thrips, scales, mealybugs, white flies, leaf miners, leaf bugs, adult Japanese beetles, lace bugs, plantbugs and tent caterpillars. Repeat treatment as necessary. Make applications for scales when crawlers settle. **CAUTION: INJURY MAY OCCUR TO CERTAIN TENDER PLANTS SUCH AS CERTAIN FERNS AND SOME SPECIES OF CRASSULA. DO NOT TREAT THESE.**

AR-267



Net Contents:

Premium Grade
Malathion WE
Emulsifiable Concentrate
50%

Product No. 6202

Active Ingredients: Malathion O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate 50%
Aromatic petroleum hydrocarbons 40%
Inert Ingredients: TOTAL 10%
TOTAL 100%

CAUTION

Keep Out Of Reach Of Children
(See side panel for other cautions)



1514 Eleventh Street
Portsmouth, Ohio 45662
Phone 614/353-1125

RTT indoor
HOUSEHOLD INSECTS: For control of Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites and Earwigs, use at the rate of 3 tablespoonfuls per quart of water (1 quart per 5 gallons). Apply as a coarse spray or with a paint brush to window sills, baseboards, drainboards, under sinks, and stoves and other areas frequented by insects.

AVOID CONTAMINATION OF FEED, FOOD UTENSILS, MILK, MILKING EQUIPMENT AND WATER.

CAUTION

Harmful if swallowed. Avoid contact with skin, wash thoroughly with warm water and soap after using. Avoid contamination of feed or food stuffs. **KEEP OUT OF REACH OF CHILDREN**

External: In case of spillage on skin, wash with warm water and soap.

Internal: Atropine is antidote. **CALL A PHYSICIAN IMMEDIATELY.**

Do not contaminate water by cleaning of equipment, or disposal of wastes.

DO NOT USE ON PETS OR WILDLIFE.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

Container Disposal

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

EPA Reg. No. 7122-42
DATE 6-27-72

Batch #


Packed by

Add to Right Panel (where indicated)

1. This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

2. APPLICATIONS WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENT

Limited to Crack and Crevice Treatment Only - Apply a small amount of material directly into cracks and crevices, such as expansion joints between equipment bases and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs, where cockroaches, waterbugs, ants, spiders, silverfish hide. Care should be taken to avoid depositing the product on to exposed surfaces or introducing the material into the air. Avoid contamination of feed or food.



FLIES: Use at the rate of 2 quarts per 25 gallon of water (1 quart to 6 gallon). For best results add one pound of sugar for each gallon of finished spray. Mix well. Spray all surfaces and places where flies and mosquitoes light or congregate. Use a wetting spray and spray to the point of run-off. Use around buildings, stockyards, dairy barns, poultry houses, garbage cans, and manure piles. Avoid contamination of feed and foodstuffs. Repeat as necessary.

FLOWERS, SHRUBS AND SHADE TREES: Use at the rate of 1 quart per 100 gallons of water (2 teaspoons per gallon). Mix well and spray plants thoroughly. This spray will kill aphids, red spider mites, bagworms, leaf-hopper, thrips, scales, mealybugs, white flies, leaf miners, leaf bugs, adult Japanese beetles, lace bugs, plantbugs and tent caterpillars. Repeat treatment as necessary. Make applications for scales when crawlers settle. **CAUTION: INJURY MAY OCCUR TO CERTAIN TENDER PLANTS SUCH AS CERTAIN FERNS AND SOME SPECIES OF CRASSULA. DO NOT TREAT THESE.**
AR-267

Emulsifiable Concentrate

50%

Product No. 6202

Active Ingredients: Malathion O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate 50%
Aromatic petroleum hydrocarbons 40%
Inert Ingredients: TOTAL 100%

CAUTION

Keep Out Of Reach Of Children

(See side panel for other cautions)

Or Chem Corp.

1514 Eleventh Street
Portsmouth, Ohio 45662
Phone 614/353-1125

OF CHILDREN

External: In case of spillage on skin, wash with warm water and soap.

Internal: Atropine is antidote. CALL A PHYSICIAN IMMEDIATELY.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

DO NOT USE ON PETS OR WILDLIFE.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Container Disposal

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

EPA Reg. No. 7122-42

DATE 6-27-72

Batch #

Packed by

Add to Right Panel (where indicated)

1. This product is toxic to fish.
Keep out of lakes, streams, or ponds.
Do not apply where runoff is likely to occur.
Do not apply when weather conditions favor drift from areas treated.
Do not contaminate water by cleaning of equipment, or disposal of wastes.
Apply this product only as specified on this label.

2. APPLICATIONS WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENT

Limited to Crack and Crevice Treatment Only - Apply a small amount of material directly into cracks and crevices, such as expansion joints between equipment bases and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs, where cockroaches, waterbugs, ants, spiders, silverfish hide. Care should be taken to avoid depositing the product on to exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

Application of this product in the food areas of food handling establishments, other than crack and crevice treatment, are not permitted.

