

## IMPORTANT

Baits should be examined regularly and replenished before completely eaten. Cereal baits prepared with Pival are resistant to insect infestation, less susceptible to mold and highly acceptable to rats and mice. However, should baits become sour, moldy or infested with insects, they must be replaced.

## CAUTION

PIVAL is an anticoagulant chemical. Therefore, PIVAL and baits prepared with PIVAL must be protected from or placed where humans, livestock, poultry, pets or wildlife will not eat them.

In case baits are eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. CALL A PHYSICIAN IMMEDIATELY.

**NOTE: FOR PHYSICIANS:** Intravenous injections of Vitamin K1 is indicated together with administration of blood transfusions.

U.S.D.A. Reg. No: 7122-84

Date

# GUARDIAN



Net Contents

## PIVAL (2%) CONC. A Rodenticide

ACTIVE INGREDIENT:		
2-Pivalyl-1,3-indandione*		2.0%
INERT INGREDIENTS:	TOTAL	98.0%
		100.0%

## CAUTION

Keep Out Of Reach Of Children  
(See side panel for additional cautions.)

*Ar Chem Corp.*

1514 Eleventh Street  
Portsmouth, Ohio 45662  
Phone 614/353-1125

## DIRECTIONS

Mix thoroughly one part of PIVAL with 76 parts of suitable bait material such as corn meal, rolled oats or other cereal grains, for 0.025% bait. A small amount of an edible oil such as mineral, corn or peanut oil may be used up to 5% to prevent dusting or to increase acceptability.

Baits should be placed in locations frequented by rats and mice and preferably where they are known to feed.

### FOR RATS:


- Four ounces of bait per station should be the **minimum** amount used.
- Bait should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.

**FOR MICE:** The amount should be reduced to a tablespoonful, but more bait placements used about every 8 to 12 feet apart. Maintain bait supply for at least 15 days.

Baits must be protected from dogs, cats, poultry, wildlife, etc. by bait boxes, cages, or other suitable means.

For complete control, baiting should be maintained continuously as long as any evidence of feeding may continue up to the fourth week. Where reinfestation is likely, permanent bait stations should be used and bait placed in them as often as necessary.

**DO NOT REUSE CONTAINER, DESTROY WHEN EMPTY.**



## IMPORTANT

Baits should be examined regularly and replenished before completely eaten. Cereal baits prepared with Pival are resistant to insect infestation, less susceptible to mold and highly acceptable to rats and mice. However, should baits become soiled, moldy or infested with insects, they must be replaced.

## CAUTION

PIVAL is an anticoagulant chemical. Therefore, PIVAL and baits prepared with PIVAL must be protected from, or placed away from, humans, livestock, poultry, cats or wildlife, which eat from them.

If your bait is eaten, give a tablespoonful of salt in a glass of warm water and report to your veterinarian. CALL A PHYSICIAN IMMEDIATELY.

**NOTE, FOR PHYSICIANS:** In severe cases, the use of Vitamin K1 is indicated, together with administration of blood transfusions.

U.S.D.A. Reg. No. 7122-84

Date:

# GUARDIAN



Net Contents

## PIVAL (2%) CONC. A Rodenticide

ACTIVE INGREDIENT	10%
INERT INGREDIENTS	90%
TOTAL	100%

## CAUTION

Keep Out Of Reach Of Children

(See side panel for additional cautions)

Or Chem Corp.

1514 Eleventh Street  
Portsmouth, Ohio 45962  
Phone 614/353-1111

## DIRECTIONS

Mix 100 parts of bait with 10 parts of PIVAL. For mice, use 1 part of PIVAL to 25 parts of bait. For rats, use 1 part of PIVAL to 25 parts of bait. A small amount of an easily eaten cereal or grain may be added to the mixture to improve its palatability.

Bait should be placed in locations frequented by rats and mice, and preferably where they are known to feed.

### FOR RATS:

1. Place 100 parts of bait per 10 parts of PIVAL. Minimum amount used.
2. Bait should be placed in areas where rats are known to feed, and where the animals are not likely to be disturbed.
3. At least 100 parts of bait should be placed in each area at least 15 days.

**FOR MICE:** The amount of bait should be increased to 25 parts of PIVAL per 100 parts of bait. Bait placements should be renewed every 15 days. Must be bait supply for at least 15 days.

Bait should be protected from access by children, cats, dogs, birds, and other animals.

For complete information on the use of PIVAL, consult the label on the container. For more information, contact your local health department or the manufacturer. PIVAL is a registered trademark of Or Chem Corp.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.