

DIRECTIONS

- Thoroughly mix one part (by weight) of rodenticide with 79 parts of dry cereal bait for 0.25 bait.
- Place bait in locations where rats and mice feed, water or travel.
- Bait placements along walls, in corners, and in concealed locations where rats and mice feed are desirable.
- Place baits in locations not accessible to children, pets, domestic animals and wildlife or in tamper-proof bait boxes.

CONTAMINATED OR SPOILED BAITS SHOULD BE REPLACED PROMPTLY.

FOR RAT CONTROL:

- Four ounces of bait per station should be the **minimum** amount used.
- An uninterrupted supply of bait should be maintained for at least 15 days.
- Bait should be placed in locations not accessible to children, pets and domestic animals or in tamper-proof bait boxes.

FOR MOUSE CONTROL:

- Tablespoon amounts of bait in each bait should be placed at 8-12 foot intervals.
- Bait should be placed in locations not accessible to children, pets and domestic animals or in tamper-proof boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.



WARFARIN 2% CONCENTRATE

RODENTICIDE

KILLS RATS AND MICE

Anti-Coagulant for Rodent Control

CAUTION

Keep Out Of Reach Of Children

(See side panel for additional cautions.)

ACTIVE INGREDIENT:

Warfarin (3-(*a*-acetyloxybenzyl)-4-hydroxycoumarin) 2.0%

INERT INGREDIENT: 98.0%

TOTAL 100.0%

Or Chem Corp.

1514 Eleventh Street
Portsmouth, Ohio 45662
Phone 614/353-1125

CAUTION

Keep away from humans, pets, domestic animals and wildlife.

If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding.

In such case, the intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of methyldihydrocoumarin.

NOTE: IF BAIT MOLDS, DECOMPOSES OR BECOMES SOUR, REPLACE WITH FRESH MATERIALS.

DO NOT REUSE CONTAINER

DESTROY WHEN EMPTY

U.S.D.A. Reg. No. 7122-85

DATE



