

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

11/ 14 7122-124 pg 17

AUG 12 1993

5446033 300/17

Archem Corporation
P.O. Box 767
1514 11th Street
Portsmouth, OH 45662

Dear Sir or Madam:

Subject: ROBAN II AG
EPA Registration No 7122-124
Your Letter Dated August 4, 1993

The revised labeling referred to above, submitted in connection with registration under FIFRA, is acceptable, providing you make the following change:

- Change the scientific name of the Polynesian rat from "(RATTUS EXVIANS)" to "(RATTUS EXULANS)".

Submit one copy of final printed labeling before you release the product for shipment with the amended labeling.

We approve your plan to sticker over the "Statement of Practical Treatment" on your existing inventory before you release the product for shipment. Note that you are still required to submit a copy of the "informational bulletin" that you must provide your distributors, as well as a list of those distributors. The Zinc Phosphide Consortium has informed you of those requirements.

This registration will be subject to cancellation in accordance with FIFRA sec. 6(e) if you do not comply with these conditions. Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

ACCEPTED
with COMMENTS
in EPA Letter Dated

AUG 12 1971

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the post-registered under EPA Reg.

7122-124

RESTRICTED

For retail sale to and use only
who are under their direct supervision
covered by the Certified Applicator

ROBAL

For control of Cotton Rats
Polynesian Rats and

Active ingredients: Zinc Phosphide
Inert ingredients:

Total

KEEP OUT OF REACH OF CHILDREN

CAUTION

Any persons applying zinc phosphide
systems such as nausea, abdominal
pain, should be seen by a physician.
IF SWALLOWED: Immediately call a physician.
Transport the patient to the nearest hospital
and administer anything by mouth or
to do so by a physician.
IF IN EYES: Flush with plenty of water
until irritation persists.
IF ON SKIN: Remove contaminated clothing
and wash with soap and water.

See side panel for

MANUFACTURER

The ArChem

1514 Elmwood
Portsmouth,
New Hampshire
Phone 614-435-1111

Net

Contents _____

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. May be fatal or harmful if swallowed. Do not breathe dust. Avoid eye or skin contact. Wash hands thoroughly after use, use rubber gloves if contact with skin may occur when placing bait. Avoid contamination of feed or foodstuffs. Avoid contact with acids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds and wildlife feeding on treated baits may be killed. Do not apply directly to lakes, streams or ponds.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original containers, in a dry place inaccessible to children, pets, and domestic animals. Do not contaminate water, feed or food by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are Acutely Hazardous. Improper disposal of excess pesticides, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed by use according to label directions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Fiber drum with liner - Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment and dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

WARRANTY

Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

AR-436

BEST AVAILABLE COPY

1000

1000

APPLICATION DIRECTIONS
SUGAR CANE

USE RESTRICTIONS: For control of POLYNESIAN, (RATTUS EXVIANS), NORWAY (R. NOB-VEGICUS), and ROOF RATS (R. RATTUS) in sugar cane fields.

BAITING SUGAR CANE FIELDS: Broadcast at the rate of 5 pounds of bait per acre. Bait may be broadcast by ground driven devices or by hand. The number of applications shall not exceed 4 per 2 year crop. 2 per 1 year crop. The total application shall not exceed 20 pounds of bait per crop cycle. Do not apply within 90 days of harvest.

DO NOT TREAT THE SAME AREA WITH ZINC PHOSPHIDE FOR 30 DAYS AFTER FIRST TREATMENT.

APPLE ORCHARDS

USE RESTRICTIONS: For control of MEADOW and PINE VOLES (MICROTUS, SPP) in apple orchard. Apply toxic bait in fall immediately after harvest and before leaf fall. Pick up drops if feasible. Do not allow animals to graze orchards or to eat drops and hay from orchards.

Where birds and other wildlife are present, handbaiting is the preferred method. Retreatment may be necessary if heavy rains follow soon after baiting or if mice move into the area later in the season.

HAND BAITING: Near the base of each infested tree, place teaspoonful quantities of bait at 2 to 4 locations, either on surface trails or at the mouth of the holes leading to underground burrow systems. Cover bait artificially (e.g. mats, boards) or by pulling overhanging grass back into place. Do not allow bait to be exposed on bare ground. Do not disturb the runway systems. Bait at the rate of 2 to 3 pounds per acre of infested trees.

TRAILBUILDING: Drop teaspoonful quantities of bait at 4 to 5 foot intervals in the artificial trail made by the machine just inside the drop line on both sides of the trees. Apply at the same rate of 2 to 3 pounds per acre.

BROADCAST BAITING: Under infested trees broadcast evenly by cyclone seeder or by hand (wear rubber gloves during hand broadcast). Concentrate in areas with the heaviest vegetative cover. Do not broadcast on bare ground or growing crops. Apply at the rate of 5 to 10 pounds per acre.

Lot No. _____ Batch No. _____ Packed By _____

ersons
uses

G
A © 1968
Rats,
BBS.

signs and
or weak-
cian, or
ter. Do
as advised
if irri-
s with

L 7122-124
7122-04-1