



**SOUTHERN STATES
FLY BAN 16% MILK MAKER (RA)**

(Contains RABON® Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.015% *
INERT INGREDIENTS	
TOTAL	99.985% ** 100.000%

* Each pound contains 68 mg RABON®.
** Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Crude Protein	(min.)	16.000%
Crude Fat	(min.)	2.500%
Crude Fiber	(max.)	9.000%
Calcium (Ca)	(min.) 0.90%	(max.) 1.400%
Phosphorus (P)	(min.)	0.700%

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain By-Products, Roughage Products, Cane Molasses, Vitamin A Palmitate (with improved stability), D-Activated Plant Sterol, Ground Limestone, Dicalcium Phosphate, Iodized Salt, Traces of: Manganous Oxide, Zinc Oxide, Iron Sulfate, Cobalt Carbonate, Copper Oxide, Calcium Iodate.

**KEEP OUT
OF REACH OF CHILDREN
CAUTION**

See back of label for additional precautionary statements.

**SEE BACK OF LABEL
FOR DIRECTIONS FOR USE**

**Manufactured By
SOUTHERN STATES COOPERATIVE, INC.
GENERAL OFFICE
RICHMOND, VIRGINIA**

FB16MM/Ra-60728-C

ACCEPTED

AUG 17 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7138-2.

**EPA Reg. No. 7138-2
EPA Est. No. 7138-OH-1**

Net Weight 100 Lbs.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When fed to cattle, RABON® Oral Larvicide passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development, RABON® Oral Larvicide helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

Fly Ban 16% Milk Maker (Ra) is designed to be fed along with your regular roughage. Do not further dilute with grain. Feed at a rate of 1.1 pounds per 100 pounds of body weight daily.

This will supply the recommended average daily intake of 70 mg. Rabon® Oral Larvicide per hundred pounds of body weight.

Start feeding early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Fly Ban 16% Milk Maker (Ra) can be fed to dry cows, replacement animals and lactating cows.

Fly Ban 16% Milk Maker (Ra) prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around dry lots, calf pens and barns to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Fly Ban 16% Milk Maker (Ra) should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.