



193

## PASTURE BALANCER CATTLE BLOCK

Aid in preventing the breeding of horn flies and face flies in the manure of treated cattle when used as directed.

### ACTIVE INGREDIENT

Phenothiazine ..... 1.00%

**INERT INGREDIENTS** .....99.00%\*

TOTAL 100.00%

\*Refers only to ingredients which are not pesticides,  
EACH POUND CONTAINS 4.57 GM PHENOTHIAZINE

ACCEPTED

FEB 22 1983

U.S. Department of Agriculture  
Federal Animal Drug Administration  
as amended, for the  
registered under  
EPA Reg. No. 7138-11

**"KEEP OUT OF REACH  
OF CHILDREN"  
CAUTION**

(See side panels for additional information.)

### PRECAUTIONARY STATEMENTS HAZARD TO DOMESTIC ANIMALS CAUTION

Keep out of reach of children. Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. Do not administer to animals showing symptoms of acute respiratory conditions. Do not use with organophosphates as phenothiazine may potentiate organophosphate toxicity. Do not administer to animals that are weak, anemic, emaciated or show any signs of severe constipation. Do not administer to pregnant animals in the last four weeks of pregnancy. Avoid exposure of recently treated animals to bright sunlight because of danger to photosensitization.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by disposal of wastes. Apply this product as specified on the label.

**MANUFACTURED BY  
SOUTHERN STATES COOPERATIVE, INC.  
GENERAL OFFICE  
RICHMOND, VIRGINIA 23260**

E.P.A. Est.No. 7138-KY-07

E.P.A. Reg. No. 7138-11

**33 1/3 LBS. NET WT.**

**3 BLOCKS EQUIVALENT TO 100 LBS.**

## **DIRECTIONS FOR USE**

It is in violation of federal law to use this product in a manner inconsistent with its labeling.

Feed this block free choice. Provide at least one block per five head of cattle and locate the blocks near the animals feeding, watering or loafing area. Do not feed salt or any other mineral supplement along with this block. Provide adequate amounts of water. Maintain adequate forage or pasture at all times to assure consistent intake.

Adult cattle must consume seven (7) ounces of Pasture Balancer Cattle Block per head per day in order for cattle to consume the 2 grams of phenothiazine intended. Younger cattle must consume approximately 9/10 ounce per 100 pounds of body weight daily to assure their receiving the 0.25 grams of phenothiazine intended per hundredweight. Maintain adequate Pasture Balancer Cattle Blocks at all times to assure continuous intake.

Cattle should be given access to Pasture Balancer Cattle Block for several weeks prior to the appearance of adult horn flies in the spring, and should be fed the supplement continuously until flies disappear in the fall. Pasture Balancer Cattle Block helps prevent the breeding of horn flies and face flies but has no effect upon existing adult fly populations. Since horn flies over-winter in the pupal state, all cattle should be sprayed when the flies first appear. Spraying kills the flies emerging from last year's pupae and will protect cattle for about two weeks until Pasture Balancer Cattle Block can take effect.

### **STORAGE AND DISPOSAL**

#### **PROHIBITIONS**

Do not contaminate water or food by storage and disposal.

#### **PESTICIDE DISPOSAL**

Feed supplement which cannot be used must be disposed of according to applicable Federal, state, or local procedures.

#### **CONTAINER DISPOSAL**

Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into feeding boxes. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

#### **STORAGE**

Do not store block next to other chemicals.

## GUARANTEED ANALYSIS

Calcium (Ca), not more than .....	4.50%
Calcium (Ca), not less than .....	3.50%
Phosphorus (P), not less than .....	1.50%
Magnesium (Mg), not less than .....	2.00%
Sulfur (S), not less than .....	0.60%
Salt (NaCl), not more than .....	18.00%
Salt (NaCl), not less than .....	16.00%
Iodine (I), not less than .....	0.0004%
Vitamin A .....	50,000 IU/lb.
Selenium (Se) (from Sodium Selenite) ...	0.0001% (1.0 p.p.m.)

## FEED INGREDIENTS

Processed Grain By-Products, Forage Products, Roughage Products, Cane Molasses, Calcium Lignin Sulfonate, Vitamin A Palmitate (with improved stability), D-Activated Animal Sterol, Magnesium Oxide, Potassium Sulfate, Magnesium Sulfate, Ground Limestone, Dicalcium Phosphate, Iodized Salt, Traces of: Manganous Oxide, Zinc Oxide, Iron Sulfate, Cobalt Carbonate, Copper Oxide, Calcium Iodate, Sodium Selenite.