NET WEIGHT 50 LBS.

LIVESTOCK FLY & WORM MINERAL MEDICATED

Aids in the control of stomachworms (<u>Haemonchus</u>, <u>Ostertagia</u> and <u>Trichostrongylus spp.</u>) nodular worms (<u>Oesophagostomum spp.</u>) large-mouth bowelworms (<u>Chabertia spp.</u>) and hookworms (<u>Bunostorum spp.</u>) in cattle when fed in accordance with directions for use. Aid in preventing in the breeding of horn flies and face flies in the manure of treated cattle when used as directed.

"KEEP OUT OF REACH OF CHILDREN" CAUTION

ACTIVE INGREDIENT

* Each pound contains 6.39 grams Phenothiazine.

** Refers only to ingredients which are not pesticides.

GUARANTEED ANALYSIS

Calcium (Ca) (min.) 17.00% (max.) 19.00%	
Salt (NaCl) (min.) 15.00% (max.) 17.00%	
Phosphorus (P) (min.) 9.00%	
Magnesium (Mg) (min.) 1.00%	
Sulfur (S) (min.) 1.00%	
Iodine (I) (min.) 0.0022%	
Vitamin A 50,000 IU/1b.	
Vitamin D	
Selenium (Se) 0.000010% (1.0 p.p.m.)	
(from Sodium Selenite)	

FEED INGREDIENTS

Processed Grain By-Products, Roughage Products (as a Vitamin Carrier), Cane Molasses, Vitamin A Palmitate (with improved stability), D-Activated Animal Sterol, Magnesium Oxide, Potassium Sulfate, Magnesium Sulfate, Ground Limestone, Dicalcium Phosphate, Iodized Salt, Traces of: Manganous Oxide, Zinc Oxide, Iron Sulfate, Cobalt Carbonate, Copper Oxide, Calcium Iodate, Sodium Selenite.

"SFF BACK PANFL FOR ADDITIONAL PRECAUTIONS"

REST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes wash with plenty of water. If irritation persists, see a physician.

Keep out of reach of children. Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. Do not administer to animals showing symptoms of acute respiratory conditions. Do not use with organophosphates as phenothiazine may notentiate organophosphate toxicity. Do not administer to animals that are weak, anemic, emaciated or show any signs of severe constination. Do not administer to pregnant animals in the last four weeks of pregnancy. Avoid exposure of recently treated animals to bright sunlight because of danger to photosensitization.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by disposal of wastes. Apply this product as specified on the label.

DIRECTIONS FOR USE

It is in violation of federal law to use this product in a manner inconsistent with its labeling.

LIVESTOCK FLY & WORM MINERAL MEDICATED must be provided to animals which are supplied with adequate roughage and/or forage at all times.

Self-feed LIVESTOCK FLY & WORM MINERAL MEDICATED as the sole source of salt. Locate product near water supply and in cattle loafing area. Maintain adequate LIVESTOCK FLY & WORM MINERAL MEDICATED at all times to assure continuous and adequate intake.

It is essential that animals fed medicated feed products have been adapted to the product. Animals that are not adapted to medicated supplementation may not consume the product at the proper level or they may overconsume the product due to inadequate prior supplementation.

Adequate LIVESTOCK FLY & WORM MINERAL MEDICATED consumption is required at all times to assure effective drug activity. It is essential that good husbandry practices be followed at all times: Medicated product feeding techniques include but are not limited to a variety of practices. Increased consumption may often be obtained by moving feeding stations into shaded or loafing areas, closer to water source, increasing feeding stations available to the animals or combinations of these practices. Decreased consumption may often be obtained by the reverse of these practices.

Medicated products should always be fed in protected feeders.

Adult cattle must consume five (5) ounces of LIVESTOCK FLY & WORM MINERAL MEDICATED per head per day in order to consume the two (2) grams of phenothiazine intended. Growing cattle must consume 0.625 ounces per 100 pounds body weight per head per day of LIVESTOCK FLY & WORM MINERAL MEDICATED to receive the 0.25 grams of phenothiazine intended.

Consult your veterinarian for assistance in diagnosis, treatment and control of parasitism.

Because this product kills only immature flies, and adulticide, in combination with good management and sanitary practices, may be necessary to control fly population.

•

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water or food by storage and disposal.

PESTICIDE DISPOSAL

Feed supplement which cannot be used must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL

Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into feeding boxes. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

STORAGE

Do not store product next to other chemicals.

Manufactured By
SOUTHERN STATES COOPERATIVE, INC.
GENERAL OFFICE

RICHMOND, VIRGINIA

E.P.A. Est.No. 7138-KY-07 NET WT. 50 LBS. E.P.A. Reg. No. 7138-

On the bottom of page 2 change "and adulticide" to "an adulticide".

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours.

George T. LaRocca Product/Manager (15) Insecticide-Rodenticide Branch Registration Division (TS-767C)