

7138-23

1/16/2015

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Rebecca M. Horton  
Consultant/Agent for  
Southern States Cooperative, Inc.  
C/o Registrations by Design, Inc.  
P.O. Box 1019  
Salem, VA 24153-1019

JAN 16 2015

Subject: Label Notifications per PRN 98-10 – Alternate Brand Name (ABN) for  
Commercial Use and Add State-specific Text for Fertilizer Components  
Product Name: Carpetmaker X-X-X with 0.25% Barricade® (ABN: Southern  
States® Professional Turf Fertilizer Active Formula Fertilizer with 0.25%  
Barricade®)  
EPA Registration Number: EPA Reg. No.: 7138-23  
Application Date: October 31, 2014  
Decision Number: 497317

Dear Ms. Horton:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, you may contact Terri Stowe at (703) 305-6117 or via email at [stowe.terri@epa.gov](mailto:stowe.terri@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn V. Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

# Carpetmaker® X-X-X with 0.25% Barricade®

[Additional Brand Name for product marketed for commercial use: Southern States® Professional Turf Fertilizer Active Formula Fertilizer with 0.25% Barricade®]

Barricade® is a Reg. Trademark of Sygenta.

For preemergence control of weed grasses and broadleaf weeds in established turf grasses as listed: in Lawns, Golf Courses, fairways, (not for putting greens), and sod farms. Use for preemergence weed control around Landscape ornamentals, including established perennials and wildflower plantings.

**NOTIFICATION**  
**JAN 16 2015**

ACTIVE INGREDIENT:	
Prodiamine***.....	0.25%
OTHER INGREDIENTS:.....99.75%	
***CAS [2909-21-2]	Total.....100.00%

### READ ENTIRE LABEL BEFORE USING

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS and DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

### FIRST AID

**IF SWALLOWED-** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES-** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING-** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED-** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. For information call 1-800-282-8007. You may also call 1-800-858-7378 for emergency medical treatment advice.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; chemical resistant gloves; and shoes plus socks and a dust mask. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Net Weight 50 Pounds

EPA Reg. No. 7138-23  
October 31, 2014

EPA Est. No. \_\_\_\_\_  
Page 1 of 12 pages

**Southern States Cooperative, Inc.**  
**P.O. Box 26234 Richmond, VA 23260 U.S.A.**

**USER SAFETY RECOMMENDATIONS**

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

**AGRICULTURAL USE REQUIREMENT**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains "specific instructions and exceptions pertaining to the" statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard for sod farms and commercial nurseries.


Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical resistant gloves, such as butyl rubber > 14 mils or natural rubber > 14 mils or Nitrile rubber > 14 mils.
- Shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The "WPS" applies when this product is used to produce agricultural plants on farms, forests, or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks, and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry.

**U.S.A. Crabgrass Seed Germination Dates**



- |                                 |                 |
|---------------------------------|-----------------|
| 1. Mid-north & northwestern     | After May 30    |
| 2. Upper/mid-states & northeast | After May 10    |
| 3. Lower mid-states             | After April 20  |
| 4. Upper southern states        | After Mar 20    |
| 5. Warm seasons states          | Jan 1 to Mar 20 |

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This product is a turf fertilizer and a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least ½ inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

## **APPLICATION DIRECTIONS**

Apply uniformly with suitable, properly calibrated, granule, application equipment.

## ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least ½ inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the U.S.A. map for approximate crabgrass seed germination dates.

## **USE PRECAUTIONS**

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. **IMPORTANT:** Do not graze or feed livestock forage or hay cut from areas treated with this product. Do not apply aerially. Failure to follow the directions for use and precautions on this label may result in poor weed control, illegal residues, damaged turf or ornamental plants.

## **Rates of Application**

This product may be applied as single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of "residual pre-emergence weed control" provided by this product is related to the rate applied.

### Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table. Maximum application rate of this product per calendar year by turfgrass species (1.)

Turf Species	Lbs. Product / Acre	Lbs. a.i. / Acre
Creeping Bentgrass	260 lbs.	0.65 lbs.
Creeping Red Fescue	300 lbs.	0.75 lbs.
Buffalograss	400 lbs.	1.00 lbs.
Kentucky Bluegrass	"	"
Perennial Ryegrass	"	"
Bermudagrass (2)	600 lbs.	1.50 lbs.
Bahiagrass	"	"
Centipedegrass	"	"
Seashore Paspalum	"	"
St. Augustinegrass	"	"
Tall Fescue	"	"
(including turf type)	"	"
Zoysia	"	"

- (1) These are the maximum rates per calendar year by species limitations.
- (2) May be used on newly-sprigged or plugged Bermuda grass at rates not to exceed 132 lbs. per Acre. Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.
- Do not apply more than 395 lbs. of this product to the same area per calendar year.
- Use the higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates as specified in the Maximum Annual Rates Table.

### Weeds Controlled

When used in accordance with this label, this product will provide control of the following weeds:

Barnyardgrass	Henbit	Purslane <i>Common</i>
Bluegrass, Annual ( <i>Poa Annua</i> )	Itchgrass	Pursely, Florida
Carpetweed	Johnsongrass ( <i>from seed</i> )	Rescuegrass (3)
Chickweed, Common	Junglerice	Shepherd's Purse (2)
Chickweed, Mouseear ( <i>from seed</i> )	Knotweed	Signalgrass, <i>Broadleaf</i>
Crabgrass ( <i>Large, Smooth</i> )	Kochia	Speedwell, <i>Persian</i>
Crowfootgrass	Lambsquarter, <i>Common</i>	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, <i>Prostrate</i>
Foxtails, <i>Annual</i>	Panicum (Texas, Fall, Browntop)	Witchgrass
Goosegrass (1)	Pigweed	Woodsorrel, Yellow ( <i>from seed</i> )

- (1) In many areas a single application of 260 to 600 lbs. per acre will control goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 260 to 400 lbs. per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates Table on this label.
- (2) For Shepherd's purse apply in late summer, fall, or winter prior to germination.
- (3) Suppression only of recuegrass.

Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

**WHEN TO APPLY AFTER OVERSEEDING TURF**

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, which ever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

**WHEN TO OVERSEED AFTER APPLICATION**

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and or reseeding results.

See adjacent map of U.S.A. for:  
**North, Transition and South, regions.**

Product Rate lbs./Acre	Months to Wait Before Overseeding			
	a.i./Acre	North	Transition	South
360	0.65	5	4	4
300	0.75	6	5	5
400	1.00	-	7	7
600	1.50	-	7	12



**LANDSCAPE ORNAMENTAL INCLUDING  
ESTABLISHED WILD FLOWER PLANTINGS**

This product may be applied for residual preemergence weed control around Ornamentals as listed.

**Use Rates**

Apply at 260 to 600 lbs. per acre (equal to 5.968 lbs. to 13.774 lbs. of this product per 1,000 sq. ft.) of the turf species surrounding landscape plants, in fall and/or spring. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate, per acre, for the turf species present. Refer to **rates chart** for species and rates. For applications over bare ground, around ornamentals, (first smooth the soil, free of clods and vegetative debris). Apply higher rates where high weed infestation is expected. Apply lower rates where low weed infestation is expected.

**Application Timing and Information**

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator. Calibrate you applicator to assure appropriate dosage as described above under "Use Rates", prior to application. Calibrate your applicator as described in the owners manual. The bulk density of this product may vary; therefore, to assure that the product is applied at the rates described, you must use a properly calibrated applicator.

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or, (in cultivated area), with a shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

**Established Landscape Ornamentals**

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

**Newly Transplanted Landscape Ornamentals**

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. Irrigate after application as above directed.

**Tolerant Ornamental Species**

This product will not harm most trees, shrubs, vines and flowers. The species listed here are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

**Application Equipment and Spreader Settings**

The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application of this product. Follow directions in the owners manual for calibrating your spreader. The settings indicated in the following table should only be used as a guide in calibrating your spreader.

<b>SPREADER SETTINGS</b>				
For Application of recommended rates of a.i. per acre				
Amt. of a.i. per acre =	0.65 lb.	0.75 lb.	1.0 lb.	1.50 lb.
50 lbs. treats sq. ft. =	8,378	7,260	5,445	3,630
<b>Spreader Settings</b>				
Scott's Evergreen 18"	# XX	# XX	# XX	# XX
Scott's Speedy Broadcast	# XX	# XX	# XX	# XX
Red Devil Broadcast	# XX	# XX	# XX	# XX
Republic ez Drop 18"	# XX	# XX	# XX	# XX
Republic ez Broadcast	# XX	# XX	# XX	# XX
Republic ez Tractor	# XX	# XX	# XX	# XX

**For Use Around landscape ornamentals  
including established perennials and wildflowers, as listed.**

<b>Scientific Name</b>	<b>Common Name</b>
<i>Abies spp.**</i>	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Achillea sp.</i>	Yarrow: King Edward
<i>Actinidia chinensis*</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African lily)
<i>Agapanthus orientalis</i>	
<i>Akebia quintata</i>	Five-Leaf of Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia sp.</i>	Aquilegia: Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemesia sp.</i>	Wormwood; Silver Mound, Castle
<i>Aster sp.</i>	Aster: Bonny Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filiz-femina</i>	Lady Fern; Fern Lady
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia sp.</i>	Fibrous Begonia: Hardy Grandis
<i>Berberis gladwynensis</i>	Barberry
<i>B. julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. thunbergii</i>	Japanese Barberry
<i>B. verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Bougainvillea spp.</i>	Bougainvillea
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Campanula carpatica</i>	Tussock Bellflower (White Clips)
<i>Campis X Tagliabuana</i>	Trumpet Creeper, Trumpet Flower, Madame Galen
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceratostigma plumbegonoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Chrysanthemum nipponicum</i>	
<i>Cleyera japonica</i>	Cleyera
<i>Citrus spp.*</i>	Citrus species*
<i>Coreopsis sp.</i>	Coreopsis (Calliopsis): Early sunrise, Moonbeam
<i>Cornus florida</i>	Flowering Dogwood
<i>C. stolonifera</i>	American Dogwood
<i>Cortaderis selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster



<i>C. buxifolius</i>	Cotoneaster
<i>C. dammeri</i>	Bearberry Cotoneaster
<i>C. microphyllus</i>	Rockspray Cotoneaster
<i>Cretaeagus spp.</i>	Hawthorne
<i>Crocasmia sp.</i>	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma sp.</i>	Cooperi Pink
<i>Delphinium sp.</i>	Larkspur: Blue Elf
<i>Dianthus deltoides</i>	Dianthus: Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea purpurea</i>	Coneflower, Purple; Magnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>E. japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>F. suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia
<i>Gaillardia sp.</i>	Gaillardia, Blanket Flower, "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine, Radicans
<i>Gaura sp.</i>	
<i>Gentiana dahurica</i>	Gentian
<i>Geranium cinereum</i>	Cranesbill
<i>Gladiolus spp.**</i>	Gladiolus species**
<i>Gypsophila repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthemum sp.</i>	Sunrose
<i>Hemerocallis sp.</i>	Daylily: Aztec Gold, Stella De Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus sp.</i>	Mallow: Disco Belle White
<i>Hibiscus, rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. Sieboldiana</i>	Hosta, "Searsucker"
<i>Houttuynia cordata var. variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangae
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly; Helleri
<i>I. opaca</i>	American Holly
<i>I. pernyi</i>	Holly
<i>I. vomatoria</i>	Yaupon Holly; Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodelsong
<i>Iris siberica</i>	Siberian Iris, Cabernet
<i>Iris spp.**</i>	Iris species**
<i>Jasminium udiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonnii
<i>J. conferta</i>	Shore Juniper, Blue Pacific
<i>J. davurca</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper

<i>Juglans sp.</i>	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender sp.</i>	Lavender, Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>L. amurense</i>	Amur Privet
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lillum sp.</i>	Lily; Jazz Big Blue Lily; Turf
<i>Liriope muscari</i>	Silver Mound; Evergreen Giant
<i>Liriope Muscari var. variegata</i>	Liripoe, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Talarian Honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum sp.</i>	Loosestrife; Modern Pink
<i>Magnolia spp.**</i>	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus sp.*</i>	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus spp.**</i>	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europea*</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterphyllus</i>	Osmanthus (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Fraser's Photinia (Redtip)
<i>Physotegia Virginiana</i>	Dragonhead, False, Vivid
<i>Picea spp.**</i>	Spruce species**(Colorado, Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. elliotii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. thunbergiana</i>	Japanese Black Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. virginiana</i>	Virginia Pine

<i>Pistachio sp.*</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus sp.*</i>	Almond, Apricot, Nectarine, Peach, Plum, Prune*
<i>Pseudotsuga menziesii**</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus Sp.</i>	Bradford Pear sp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus spp.</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorn
<i>Rhododendron</i> (including Azalea)	Coral Bells, Delaware Valley White, Flame Creeper, Formosa, Girard Crimson, George L. Tabor, Hino-crimson, PJM, Roseum Elegans, Wakeiebisu, White Gumbo
<i>Rosa banksiae*</i>	Lady Bank's Rose
<i>Rosmarinus officinalis*</i>	Rosemary*
<i>Rudbeckia sp.</i>	Black-Eyed Susan; Goldstrum
<i>Santolina, virens</i>	Santolina
<i>Saxifraga sp.</i>	Saxifrage; Purple Dome
<i>Scabiosa sp.</i>	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum caudicicola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea Bumaida</i>	Spirea; Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidate</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium sp.</i>	Germander
<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa spp.</i>	Tulip species
<i>Veronica sp.</i>	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowwood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca minor</i>	Common Periwinkle, Vinca
<i>Vitis Sp.*</i>	Grape*

<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filamentosa</i>	Adam's Needle, Yucca

**\*Do not use on food production trees, vines, or plants.**  
**\*\*Not for use on container grown plants.**

**STORAGE AND DISPOSAL**

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.  
**PESTICIDE STORAGE:** Store in original labeled container, in a dry, cool area out of reach of children and pets. Partially used bags should be sealed by folding down the top and securing.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this bag. Completely empty bag into application equipment. Offer for recycling, if available. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned stay out of smoke.

**GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N).....variable%  
 Ammoniacal Nitrogen.....variable%  
 Water Soluble Nitrogen.....variable%  
 (and/or Urea Nitrogen)  
 Water Insoluble Nitrogen.....variable%  
 Available Phosphate (P<sub>2</sub>O<sub>5</sub>).....variable%  
 Soluble Potash (K<sub>2</sub>O).....variable%

Derived from XXX.  
 Chlorine not more than .....variable%

[Optional State-specific text corresponding to fertilizer components:

FLORIDA APPLICATIONS: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N< P< K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://www.edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>

FLORIDA APPLICATIONS: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except for golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at <http://www.dep.state.fl.us/Central/Home/Meetings/Training/FLGreen/BMP Book final.pdf>

**If fertilizer formula contains P and is sold into MD or VA – the applicable following text may be featured:**

MD: This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) reestablishing or repairing a turf area.

MD: (IF PHOS >5% & not a "starter"): front panel, 3/4 inch font: NOT FOR USE ON ESTABLISHED LAWNS OR GRASS IN MD

VA: In Virginia this lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.]

**CONDITIONS OF SALE AND WARRANTY**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Southern States Cooperative, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, there are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Southern States Cooperative, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Southern States Cooperative, Inc. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

**LIMITATION OF WARRANTIES:** To the extent consistent with applicable law the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Southern States Cooperative, Inc. option, the replacement of the product.

[Lot no. to be added at production.]