

Jolt

SHOCK TREATMENT


ACTIVE INGREDIENTS:
TRICHLORO-S-TRIAZINETRIONE 100%

EPA Reg. No. 7152-9

**DANGER: KEEP OUT OF REACH
OF CHILDREN**

See side panel for antidote and other precautions.

NET WT. 2 lbs.



**SEABOARD
INDUSTRIES**
HAWTHORNE, NEW JERSEY

DIRECTIONS:

For newly-filled pools.

To establish a chlorine residual, add 1 pound of Jolt for every 8,000 gallons of water. Repeat dosage until a chlorine residual of 1.0 to 1.5 ppm is obtained as determined by the use of a chlorine test kit.

Thereafter, repeat the above dosage daily or as needed to maintain 1.0 to 1.5 ppm chlorine residual at all times.

Test water frequently with a chlorine test kit. Keep the pH of the pool water in the 7.2 to 7.6 range.

Danger: Keep out of reach of children. May be fatal if swallowed. Strong oxidizing material. May cause eye damage. Protect eyes when handling. Dust may be harmful if inhaled. Avoid contact with skin, eyes and clothing. May cause burns to broken skin. In case of contact, immediately flush skin and eyes with cold water for at least 15 minutes. For eyes, get medical attention - avoid breathing dust. Contact with organic matter may cause fire. Moisture contamination may liberate chlorine-containing gases. Keep away from open flames. Keep container closed and in a cool place when not in use. Rinse container thoroughly when discarding.

ANTIDOTE

EXTERNAL: Immediately flush with plenty of cold water.

INTERNAL: Feed bread soaked in milk followed by olive oil or cooking oil. Call a physician.

HOW TO USE CHLORINE SANITIZER

To start your pool: 1. Backwash filter. 2. Using your pool test kit adjust pH of pool water to 7.2—7.6. Add soda ash or sodium bicarbonate to correct an acid condition (less than 7.2). Add sodium bisulfate if the pH is greater than 7.6. Maintain pH range of 7.2—7.6 for a minimum of six hours before the addition of Chlorine Sanitizer and during the entire season. 3. Use two cups of Chlorine Sanitizer per 5,000 gallons of water.

After the proper pH range (7.2-7.6) and chlorine residuals (1.0-1.5) have been established, it is beneficial to add a chlorine stabilizer (cyanuric acid) to the pool to prevent extreme fluctuations in the chlorine level. Cyanuric acid is available at most pool dealers and should be added in accordance to the instruction on the package for maximum stabilization of the pool water.

HOW TO MAINTAIN THE PROPER CHLORINE RESIDUAL

The chlorine residual should be maintained between 1.0 and 1.5 ppm available chlorine as determined by the chlorine test kit. Under normal conditions, this residual can be maintained by the daily addition of about 1/2 cup (2.7 ounces) of Chlorine Sanitizer for each 5,000 gallons of pool water — if the pool water is properly stabilized with 30—50 ppm of cyanuric acid pool stabilizer. Severe conditions, such as high temperature or continued use by many people, will require higher daily dosage, whereas smaller doses may suffice in spring and fall or when the pool is not being subjected to normal bathing loads.

The pool water should be super-chlorinated about once each week. Five times the normal daily addition, or a minimum of one cup (eight ounces) per 5,000 gallons of pool water should be added in the evening when the pool will not be used for a minimum of 12 hours. Test the chlorine residual and if above 2.0 ppm do not allow swimming until residual has decreased.

SEABOARD sea/chlor

**DRY CHLORINE SANITIZER
FOR ALL SWIMMING POOLS**

- Easy to Use • Just Pour into Pool
- Clears Water Instantly • Completely Soluble
- Highly Effective Bactericide and Disinfectant

Contains 35 % available chlorine.


Active Ingredient, lithium hypochlorite 29%

Inert Ingredients 71%

WARNING: Keep out of children's reach
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS.

EPA REG. NO. 7152-15

Net.Wt. 4 1/2 Lbs.



**SEABOARD
INDUSTRIES**
HAWTHORNE, NEW JERSEY

ACCEPTED

UNIVERSITY
POOL
7/57



DIRECTIONS:

For newly-filled pools.
To establish a chlorine residual, add 1 pound of Jolt for every 8,000 gallons of water. Repeat dosage until a chlorine residual of 1.0 to 1.5 ppm is obtained as determined by the use of a chlorine test kit.
Thereafter, repeat the above dosage daily or as needed to maintain 1.0 to 1.5 ppm chlorine residual at all times.
Test water frequently with a chlorine test kit. Keep the pH of the pool water in the 7.2 to 7.6 range.

Danger: Keep out of reach of children. May be fatal if swallowed. Strong oxidizing material. May cause eye damage. Protect eyes when handling. Dust may be harmful if inhaled. Avoid contact with skin, eyes and clothing. May cause burns to broken skin. In case of contact, immediately flush skin and eyes with cold water for at least 15 minutes. For eyes, get medical attention. Avoid breathing dust. Contact with organic matter may cause fire. Moisture contamination may liberate chlorine-containing gases. Keep away from open flames. Keep container closed and in a cool place when not in use. Rinse container thoroughly when discarding.

ANTIDOTE

EXTERNAL Immediately flush with plenty of cold water.
INTERNAL Feed bread soaked in milk followed by olive oil or cooking oil. Call a physician.