

NET CONTENTS

POUND

WARNING (18p) Keep out of Reach of Children (12p.)



PRECAUCION Manténgase Lejos del Alcance de los niños

CHEMPAR MALATHION 25W

KILLS FLIES, CERTAIN ORCHARD, GARDEN, AND LIVESTOCK INSECTS

ACTIVE INGREDIENT: Malathion 25.0% INERT INGREDIENTS 75.0% 100.0%

INGREDIENTE ACTIVOS Malathion 25.0% INGREDIENTES INERTES 75.0% 100.0%

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY HOW TO PREPARE FLY CONTROL SPRAYS AND BAITS

ANTES DE USAR LEA LAS PRECAUCIONES CUIDADOSAMENTE

Table with columns for STRAIGHT SPRAYS and BAIT SPRAYS, detailing amounts of Malathion 25-W and other ingredients like Sugar and Molasses.

Table with columns for ROCIO REGULAR and BAIT SPRAYS, detailing amounts of Malathion 25-W and other ingredients like Sugar and Molasses.

\*Do not use Molasses treated with preservative.

\*Do not use Molasses treated with preservative.

DIRECTIONS

FLY CONTROL: For the reduction of flies in and around buildings which house domestic animals, around yards and homes, apply spray or bait prepared according to table on front of label.

PLANT AND ANIMAL USES

THOROUGH COVERAGE WITH SPRAY IS ESSENTIAL FOR GOOD CONTROL. ALL RATES OF APPLICATION ARE FOR USE IN 100 GALLONS OF WATER

Small Volume Rate:

5 level teaspoons per gallon = 2 1/2 lbs./100 gal. water

VEGETABLES: (Apply 100 gal. of spray per acre or 1 gal. per 435 square feet.) To control spider mites and aphids on Tomatoes and Eggplants; aphids on Peppers, Lettuce and Brussel Sprouts; aphids and leafhoppers on Potatoes, apply 3 1/2 pounds.

FRUIT: Spray with 2 1/2 pounds for control of rose apple aphids, forbes scale, and mealybug on Apples; mealybug on Pears; leafhoppers, black-headed fireworms, spittlebug nymphs, and cranberry fruitworm on Cranberries; aphids and spider mites on Strawberries.

Dilute 8 pounds for control of codling moth, plum curculio and red-banded leaf-roller on Pears and Apples, also fruit tree leafroller on Pears; Oriental fruit moth, plum curculio on Peaches. Apply 1-1 1/2 pounds for leafhopper control on Grapes. Use 1 pound for control of mealy plum aphids on Plums and Prunes.

\*Make at least two applications 10-12 days apart in summer months.

Malathion 25-W may injure Bosc Pears in the Northeast or some varieties of Sweet Cherries, especially in the Northwest.

On all applications repeat as required for continued control. Do not apply within 3 days of harvest on Apples, Pears, Cherries, Cranberries, Grapes, Plums, Prunes, Strawberries, 7 days on Peaches. May cause injury to Ribier Grapes.

ANIMALS: To control lice on Beef Cattle and Hogs use 16 pounds per 100 gallons of water, for control of horn flies on Beef Cattle use 16 to 24 pounds, for ticks on Beef Cattle use 16 to 22 pounds. Apply a complete coverage spray. Apply also to hog pens and litter.

CAUTION: Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of foodstuffs, food containers, feed and watering troughs.

NOTICE: The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed, available research indicates that this product will give satisfactory results and that the residue will be within official tolerances.

INSTRUCCIONES PARA CONTROLAR LAS MOSCAS:

Para reducir el numero de moscas en los edificios y sitios circundantes donde se alojan los animales domesticos, alrededor de los solares y las casas, apliquese un rocío o un cebo preparado de acuerdo con el índice al frente de la etiqueta.

PARA USO EN LAS PLANTAS Y LOS ANIMALES

PARA BUEN CONTROL ES ESENCIAL ESTAR BIEN ROCIADO TODAS LAS PORCIONES DE APLICACION SON PARA USARSE EN 100 GALONES DE AGUA

PROPORCION PARA PEQUEÑO VOLUMEN:

5 cucharaditas rasas por galon = 2 1/2 libras/100 galones de agua. VEGETALES: (Apliquense 100 galones de rocío por hectarea o un galon por cada 435 pies cuadrados).

FRUTA: Apliquense 2 1/2 libras de rocío para combatir los pulgones rosados de los manzanos, "forbes scale" y cochinilla harinosa de los manzanos, cochinilla harinosa en los perales, saltarillas, el gusano cocuyo de cabeza negra, las crisálidas del gusano medidor y el gusano de los arandanos, tambien como los pulgones y los acaros en los fresales.

Diluyanse 3 libras para controlar el gusano de la manzana, el gusano del ciruelo, y las chinches de manchas rojas en las peras y manzanas; el gusano oriental, el gusano del ciruelo en los melocotones o duraznos.

\*Házanse por lo menos dos aplicaciones con 10 a 12 días de diferencia durante los meses de verano.

El Malathion 25-W puede dañar las peras Bosc en el noreste o algunas variedades de cerezas dulces, especialmente en el noroeste.

ANIMALES: Para controlar los piojos en el ganado vacuno y los cordos, usense 16 libras por cada 100 galones de agua; para controlar las moscas en los cuernos del ganado vacuno usense de 16 a 24 libras; para las garrapatas en el ganado vacuno usense de 16 a 22 libras.

ADVERTENCIA: Es nocivo si se traga. Evítase a respirar prolongadamente el polvo o la niebla de la rociadura. Evítase el contacto y olfateo o repetido con la piel lavase bien despues de usarlo.

Do NOT Store with Foodstuffs

Manufactured by

CHEMPAR CHEMICAL CO., INC.

260 MADISON AVENUE NEW YORK 16 N. Y. U.S.A.

15