

181

**PRECAUTIONARY STATEMENTS:
HAZARDS TO HUMANS &
DOMESTIC ANIMALS**

CAUTION

Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes,
Streams or Ponds.

73 20

STORAGE AND DISPOSAL

**DO NOT CONTAMINATE WATER, FOOD OR FEED
BY STORAGE OR DISPOSAL**

STORAGE: Store in original container in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

191

Warranty: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



**READY-TO-USE
RAT AND MOUSE BAIT**

Active Ingredient:

2-[(p-chlorophenyl) phenylacetyl]

1,3 indandione*.....0.005%

Inert Ingredients.....99.995%

Total.....100.000%

*Chlorophacinone-Liphadione

**KEEP OUT OF REACH OF
CHILDREN
CAUTION**

See left panel for additional precautionary
statements

Manufactured by:

LiphaTech, Inc.

3101 W. Custer Avenue

Milwaukee, WI 53209 U.S.A.

EPA Reg. No. 7173-80 EPA Est. No. 7173-WI-01

NET CONTENTS: 40 LBS (18.14 Kg)

ACCEPTED
APR 26 1981
U.S. DEPARTMENT OF AGRICULTURE
NATIONAL CENTER FOR ENVIRONMENTAL HEALTH
EPA Reg. No. 7173-80

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS - For control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings, and similar manmade structures.

Treated baits must be placed in tamper resistant bait boxes or in locations not accessible to children, pets, domestic animals, or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS -

Determine areas where rats/mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.