

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
WARNING

INHALATION HAZARD

Tracking powder dust is a lung irritant. If inhaled, it can cause throat irritation, difficulty breathing and possibly fatal respiratory disease. Domestic animals exposed to tracking powder may reduce their climbing ability or may cause bleeding. In such cases, intravenous and oral administration of Vitamin K₁ combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of warfarin or dicumarol. Call a physician immediately.

Do not contaminate feed or food stuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife. Keep out of lakes, streams, or ponds.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed stuffs, storage or disposal.

PESTICIDE DISPOSAL

Pesticide spray mixture or residue that cannot be cleaned according to label instructions must be disposed of according to Federal, State or Local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Take rinses (or equivalent) and water for recycling or reconditioning and dispose in sanitary landfill or by incineration, per state, county, city and local authority.

NOTICE: Batteries may contain mercury. Do not dispose of batteries in regular trash. Do not burn batteries. Do not allow batteries to come in contact with water. Do not handle batteries when hands are wet and do not touch metal parts of batteries.

RESTRICTED USE PESTICIDE

For rate see label. Only by Certified Applicators or persons under their direct supervision for those uses covered by the Certified Applicators Certificate.

ROZOL

**TRACKING POWDER
FOR MICE AND RATS
FOR INDOOR USE ONLY**

CONTENTS:	LBS.
Active ingredient 2-[p-chlorophenyl] phenylketone [*]	0.0
1,2-indandione [*]	99.8
Inert ingredients	0.0
	100.0

*Ciba-Geigy Corporation
Patents in the U.S.A.

**WARNING
KEEP OUT OF REACH
OF CHILDREN
SEE SIDE PANELS FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.**

**MANUFACTURED BY
CHEMPAR CHEMICAL CO., INC.**

60 EAST 42nd STREET
NEW YORK, N.Y. 10155
(212) 687-3990

EPA REG. NO. 2173-113
EPA EST. NO. 3532-NY-01

ACCEPTED

JUN 19 1981

100-100-100-100
EPA Reg. No. 2173-113

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: Use against the Norway, rat, roof rat and house mouse in and around residential and agricultural buildings and similar man-made structures. ROZOL Tracking Powder may also be dusted into rat burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures. Tracking powder must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place powder in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not place near ventilation duct openings.

SELECTION OF TREATMENT AREAS: Determine dry areas where rats and mice will most likely pick up the powder on their feet or fur and ingest it during grooming. Generally, these areas are along walls, by gnawed openings and burrows, in corners and concealed places, in spaces between floors and walls, or in locations where rodents or their signs have been observed. Remove goods piled directly on the floor and place on shelves. Use boxes or other obstacles to force rodents to travel through constricted areas. Give special attention to the climbing ability of roof rats. For this species, use suitable cardboard tubes or wooden tunnels securely attached to rafters or other horizontal surfaces where rats will pass. Employ tubes large enough to minimize spillage of powder through the ends.

APPLICATION DIRECTIONS: For house mice use at the rate of 1 oz. powder per 2.5 sq ft of runway area. For Norway and Roof rats use at the rate of 2 oz. powder for 2.5 sq ft of runway area. Apply the powder into the rodent burrow or holes or within walls with a hand bulb or similar duster. Do not use power dusting devices. Sprinkle the powder in patches in such a manner as to expose the rodent to it. Patch size may be (but not limited to) 6" x 12" and should be adapted to each situation. For burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures, place about 5 grams of tracking powder in each burrow with foot-dump duster (about 15 pumps). Close burrows with soil, loose leaves, or paper. Retreat if burrows are reopened.

Maintain powder in treated areas for at least 20 days. Collect and dispose of all dead animals and used powder promptly.

Repeat treatment when infestation recurs. Where a continuous source of infestation is present, establish permanent tracking powder stations and replenish the powder as needed.

BEST DOCUMENT AVAILABLE