

ACCEPTED

JAN 15 1981

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, this pesticide
registration number
EPA Reg. No. 7173-171

maki[®]



RAT AND MOUSE MEAL BAIT

FOR INDOOR USE ONLY

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins

CONTENTS:	LBS.
Active ingredient:	
3-[3-(4-Bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	0.005%
Inert ingredients	99.995%
	100.000%

*Bromadiolone-Supercard
Patented in the U.S.A.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wash arms, and face with soap and water after applying and before eating and smoking. Exposure to Maki during pregnancy should be avoided. If swallowed, call a physician at once.

See left panel for additional precautionary statements.

CHEMPAR CHEMICAL CO., INC.

60 EAST 42nd STREET
NEW YORK, N.Y. 10165

EPA REG. NO. 7173-171
EPA EST. NO. 45221-CN 01

KEEP AWAY FROM FOODSTUFFS

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND
DOMESTIC ANIMALS
CAUTION**

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of the blood and cause bleeding.

Do not get in eyes, or skin, or on clothing. Wash arms, hands, and face with soap and water after applying and before eating and smoking. Exposure to Maki during pregnancy should be avoided. If swallowed, call a physician at once.

NOTE TO PHYSICIAN-If ingested, administer Vitamin K, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary, based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. OPEN BURNING AND DUMPING IS PROHIBITED. DO NOT REUSE EMPTY CONTAINER.

PESTICIDE DISPOSAL: Pesticide that cannot be used, should be disposed of in a landfill approved for pesticides or buried in a safe place, away from water supplies.

CONTAINER DISPOSAL: Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS For control of Norway rats, Roof rats and House mice in homes, industrial buildings and commercial buildings. Maki may also be used inside transport vehicles (ships, trains, aircraft) and inside related port or terminal buildings. For indoor use only. Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

**APPLICATION DIRECTIONS:
NORWAY AND ROOF RATS**

Apply 4 to 16 ounces of bait (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

HOUSE MICE

Apply 1 to 1.5 ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

RATS & MICE

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.