

172

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND
DOMESTIC ANIMALS
CAUTION**

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of the blood and cause bleeding.

Do not get in eyes, on skin, or on clothing. Wash arms, hands, and face with soap and water after applying and before eating and smoking. If swallowed, call a physician at once.

NOTE TO PHYSICIAN:

If ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL

Storage:

Store only in original container, in a dry place inaccessible to children and pets.

Disposal:

Do not reuse empty container. Securely wrap in newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

maki®

**RAT AND MOUSE MEAL BAIT (PACKS)
FOR INDOOR AND OUTDOOR
RODENT CONTROL**

**KILLS NORWAY RATS, ROOF RATS, HOUSE MICE
AND WARFARIN RESISTANT NORWAY RATS**

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

CONTENTS: 1½ oz (43gm)

Active Ingredient:

3-[3-(4'-Bromo[1,1'-biphenyl]-4-yl)-

3-hydroxy-1-phenylpropyl]-4-hydroxy-

2H-1-benzopyran-2-one

0.005%

Inert Ingredients

99.995%

100.000%

*Bromadolone-Supercid

Patented in the U.S.A.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wash arms, and face with soap and water after applying and before eating and smoking. If swallowed, call a physician at once.

See left panel for additional precautionary statements.

Chempar® Products

division of

LIPHA CHEMICALS, INC.

680 MADISON AVENUE

NEW YORK, NY 10021

EPA Reg. No. 7173-171

EPA Est. No. 7173-NY-01

KEEP AWAY FROM FOODSTUFFS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

For control of Norway rats, Roof rats and House mice.

Urban Areas: May be used in and around the periphery of homes, industrial, commercial and public buildings. Maki may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. May also be used in alleys. Do not use in sewers.

Nonurban Areas: May be used inside homes and agricultural buildings.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS:

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS:

NORWAY AND ROOF RATS:

Place at least 3 to 10 packs of bait (usually 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

HOUSE MICE:

Open the pack and place tablespoon amount (¼ to ½ oz.) of bait at intervals of 8 to 12 feet per placement. One pack may be placed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

RATS & MICE:

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

BEST COPY AVAILABLE

7173-171

PC114

182

8/2

ACCEPTED
JUN 11 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 7173-171

BEST COPY AVAILABLE

2000
04-88

13

2