

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
WARNING**

Harmful if swallowed

Do not inhale powder. Wear dust respirator. This product can be absorbed through the skin so it should be handled with gloves. Keep away from humans, domestic animals or pets; this material may reduce the clotting ability of the blood and cause bleeding. In such cases intravenous and oral administration of Vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Do not contaminate feed or foodstuffs

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife. Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary landfill or by incineration if permitted by State and local authorities.

NOTICE: Seller, maker, no warranty expressed or implied, concerning the use of this product or than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.



RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators Certificate.

ROZOL

**BLUE TRACKING POWDER
FOR MICE AND RATS
FOR INDOOR USE ONLY**

CONTENTS:

LBS.

Active ingredient

2-[p-chlorophenyl] phenylacetyl]-

1,3-indandione*

Inert ingredients

0.2%
99.8%

*Chlorophacchon-Lipadione
Patents in the U.S.A.

**WARNING
KEEP OUT OF REACH
OF CHILDREN**

SEE SIDE PANELS FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

MANUFACTURED BY
CHEMPAR CHEMICAL CO., INC.

60 EAST 42nd STREET
NEW YORK, N.Y. 10165
(212) 687-3990

EPA REG. NO. 7173-172
EPA EST. NO. 36302-NY-01

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: Use against the Norway rat, roof rat, and house mouse inside of homes, industrial and agricultural buildings, and similar man-made structures. ROZOL Tracking Powder may also be dusted into rat burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures. Tracking powder must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place powder in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not place near ventilation duct openings.

SELECTION OF TREATMENT AREAS: Determine dry areas where rats and mice will most likely pick up the powder on their feet or fur and ingest it during grooming. Generally, these areas are along walls, by gnawed openings and burrows in corners and concealed places, in spaces between floors and walls or in locations where rodents or their signs have been observed. Remove goods piled directly on the floor and place on skids. Use boxes or other obstacles to force rodents to travel through constricted areas. Give special attention to the climbing ability of roof rats. For this species, use suitable cardboard tubes or wooden tunnels securely attached to rafters or other horizontal surfaces where rats will pass. Employ tubes long enough to minimize spillage of powder through the ends.

APPLICATION DIRECTIONS: For house mice use at the rate of 1 oz. powder per 2.5 sq. ft. of runway area. For Norway and Roof rats use at the rate of 2 oz. powder for 2.5 sq. ft. of runway area. Apply the powder into the rodent burrow or holes or within walls with a hand bulb or similar duster. Do not use power dusting devices. Sprinkle the powder in patches in such a manner as to expose the rodent to it. Patch size may be (but not limited to) 6"x12" and should be adapted to each situation. For burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures, place about 5 grams of tracking powder in each burrow with foot-dump duster (about 15 pumps). Close burrows with soil, loose leaves or paper. Retreat if burrows are reopened.

Maintain powder in treated areas for at least 20 days. Collect and dispose of all dead animals and used powder promptly.

Repeat treatment when infestation recurs. Where a continuous source of infestation is present, establish permanent tracking powder stations and replenish the powder as needed.

ACCEPTED
 JUN 19 1981
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7173-172