

Reg # 7173 - 185

PM - 14

191

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Keep away from humans, domestic animals and pets. Keep away from feed or foodstuffs. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding.

This product should be handled with gloves. Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking. If swallowed, call a physician immediately.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

STORAGE: Store in original container in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse or equivalent. The offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burn 1, stay out of smoke.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the product label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ROZOL

LAQ-BERRY

RAT AND MOUSE BAIT

FOR INDOOR AND OUTDOOR RODENT CONTROL.

KILLS NORWAY RATS, ROOF RATS AND HOUSE MICE.

ACTIVE INGREDIENT:

| | |
|---------------------------------------------------|-----------------|
| 2-[(p-Chlorophenyl) phenylacetyl]-1,3-indandione* | 0.005% |
| INERT INGREDIENTS..... | 99.995% |
| TOTAL: | 100.000% |

*Chlorophacinone-Liphadione

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED BY:

LiphaTech, Inc.
3101 West Custer Ave.
Milwaukee, WI 53209

EPA Reg. No. 7173-185
EPA Est. No. 7173-WI-001

NET WEIGHT: 45 lbs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If a bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.


SELECTION OF TREATMENT AREA: Determine areas where rats and or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8- to 12-foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.



BEST AVAILABLE COPY