

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

APR 30 2003

Mr. Tom Schmit
Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209

Dear Ms. Schmit:

Subject: Addition of Packaging Sizes on Labeling
Maki Rat and Mouse Bait Packs (Pellets)
EPA Registration No. 7173-188
Your Submission Dated March 5, 2001

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided you make the following changes before you release the product for shipment bearing the amended labeling:

1. For both the outer container and place pack labels, change the modified single-feeding claim as follows: "Norway rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins."
2. In the second sentence of the "Urban areas:" paragraph, change "in and around transport vehicles" to "in transport vehicles."
3. In the third sentence of the "Rats and Mice:" paragraph, change "To prevent reinfestation" to "To discourage reinfestation."
4. Insert the following as the second sentence of the "Application Directions" for "Rats:": "Space placements at intervals of 15 to 30 feet in infested areas."

A stamped copy is enclosed for your records. If you have any questions, please contact me at (703) 308-6249 or hebert.john@epa.gov.

Regards,

A handwritten signature in black ink, appearing to read "John Hebert", written over a large, stylized loop.

John Hebert
Insecticide-Rodenticide Branch
Registration Division (7505C)

Outer Container Label

Rats: Place at least 3 to 10 packs per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

Mice: Use one pack per bait placement. Space placements at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

Rats and Mice: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding. Do not get in eyes, on skin, or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking. If swallowed, call a physician at once.

Note to Physician: If ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL:
Do not contaminate water, food or feed by storage or disposal. Store in original container in locked storage area.

Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Securely wrap excess, unused or spoiled bait in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse pail. Rinse thoroughly before discarding in trash.

KEEP AWAY FROM FOODSTUFFS

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 7173-188
EPA Est. No. 7173-WI-1

MAKI

RAT AND MOUSE BAIT PACKS (PELLETS)

FOR INDOOR AND OUTDOOR
RODENT CONTROL

KILLS NORWAY RATS,
ROOF RATS, HOUSE MICE AND
WARFARIN-RESISTANT NORWAY RATS

NORWAY RATS AND HOUSE MICE MAY
CONSUME A LETHAL DOSE IN ONE
FEEDING WITH FIRST DEAD RODENTS
APPEARING FOUR OR FIVE DAYS
AFTER TREATMENT BEGINS

Active Ingredient:
*3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one 0.005%
Inert Ingredients 99.995%
Total 100.000%

*Bromadiolone-Supercid Patented in the U.S.A.

KEEP OUT OF REACH OF CHILDREN
CAUTION
May be harmful or fatal if swallowed. See side panel for additional precautionary statements.

Manufactured By:
LIPHATECH, INC.
3101 W. Custer Avenue
Milwaukee, WI 53209

NET WEIGHT:

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway Rats, Roof Rats and House Mice.

Urban areas: This product may be used in and around the periphery of homes, industrial, commercial and public buildings. This product may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. This product may be used in alleys.

Non-urban areas: This product may be used inside of homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:
(Qualifier statement for 3 x 1.5 oz package only): The bait packs in this package contain enough bait (4.5 oz.) to kill one to four rats.
(Qualifier statement for 5 x 1.5 oz package only): The bait packs in this package contain enough bait (7.5 oz.) to kill two to seven rats.
(Qualifier statement for 6 x 1.5 oz package only): The bait packs in this package contain enough bait (9 oz.) to kill three to nine rats.
(Qualifier statement for 8 x 1.5 oz package only): The bait packs in this package contain enough bait (12 oz.) to kill three to twelve rats.

Place Pack Label

ACCEPTED
with COMMENTS
In EPA Letter Dated:
APR 30 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7173-188.

793-288

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

KEEP AWAY FROM FOODSTUFFS

STORAGE AND DISPOSAL:
Do not contaminate water, food or feed by storage or disposal. Store in original container in locked storage area.

Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Securely wrap excess, unused or spoiled bait in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse wrapper. Securely wrap original container in several layers of newspaper and discard in trash.

MAKI

RAT AND MOUSE BAIT PACKS (PELLETS)

FOR INDOOR AND OUTDOOR
RODENT CONTROL

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE
AND WARFARIN-RESISTANT NORWAY RATS

NORWAY RATS AND HOUSE MICE MAY
CONSUME A LETHAL DOSE IN ONE FEEDING
WITH FIRST DEAD RODENTS APPEARING
FOUR OR FIVE DAYS AFTER TREATMENT BEGINS

Active Ingredient:
*3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one 0.005%
Inert Ingredients 99.995%
Total 100.000%

*Bromadiolone-Supercid Patented in the U.S.A.

EPA Reg. No. 7173-188 EPA Est. No. 7173-WI-1

KEEP OUT OF REACH OF CHILDREN
CAUTION: May be harmful or fatal if swallowed.
See side panel for additional precautionary statements.

LIPHATECH, INC.
3101 West Custer Ave.
Milwaukee, WI 53209 USA
(414) 351-1476

NET WEIGHT: 1.5 oz (43 grams)

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label on the outer package before using this product. It is illegal to sell these placepacks individually.

PRECAUTIONARY STATEMENTS: Hazard to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding. Do not get in eyes, on skin, or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking. If swallowed, call a physician at once.

Note to Physician: If ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.