

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION**

Keep away from humans, domestic animals and pets. Do not breathe fumes or dust. May be fatal if swallowed. Do not allow contact with skin. Wash hands after use. Do not use on or near feed or foodstuffs. Keep away from children. Wear gloves and protective clothing when using bait. Do not use or store in house.

**ENVIRONMENTAL HAZARDS**

This product is toxic to wildlife and fish. Birds and other wildlife feeding on treated bait may be killed. Do not apply to open ground, roads, and areas in which no vegetation occurs. Pick up and properly dispose of all spilled toxic bait. Keep out of lakes, streams, ponds, and aquatic systems. Do not apply where runoff from this product will enter an aquatic system. Check for and dispose of excess spoiled bait and carcasses daily until the bait is expended or removed.

**ENDANGERED SPECIES  
CONSIDERATIONS**

Before field use of this product, consult Federal, State, and Local game authorities to determine if the following wildlife are found in the area. If so, observe the following:

- Whooping Crane (*Grus americana*)  
Do not use this product 30 days prior to arrival and 30 days after the whooping crane leaves known crane habitat.
- Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*)  
Do not use this product in the presently occupied range of the Attwater greater prairie chicken in the following Texas counties, Arkansas, Austin, Brazoria, Colorado, Dewitt, Fort Bend, Galveston, Galad, Harris, Refugio, Victoria, Walker, and Wharton.
- Yellow-Shouldered Black Bird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba nornata wetmorei*)  
This product must not be used in the following areas except in tamper proof bait boxes (within 10 kilometers of central Acuirre, Lago Cidra, Ceiba, San German, 15 Kilometers of La Esperanza south of Highway 2 from city of Mayaguez to the city of Ponce and all of Mona Island, Puerto Rico).
- Utah Prairie Dog (*Cynomys parvidens*)  
Do not use this product in the range of the Utah Prairie Dog (Utah).
- Salt Marsh Noddy (*Rostronodops raviventris*)  
Do not use this product within 1/2 mile of salt marsh vegetation and/or brackish water wetlands which are located (1) near or adjacent to Salt Pable Bay and San Francisco Bay or (2) in the Sacramento River below or adjacent to the confluence of the Sacramento River and the San Jaquin River (California).

**ENDANGERED SPECIES CONSIDERATIONS (cont)**

- Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*)  
Do not use this product within 1/2 mile of Baywood Park which is located on Morro Bay (California).
- Aleutian Canada Goose (*Branta canadensis leucoparia*)  
Do not use this product from October to March in the Federally closed goose hunting areas in the state of California. See current Duck & Goose Hunting game laws for current maps.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This bait must not be applied on roads, near residential areas, or over water. Do not broadcast over growing crops other than sugarcane. Apply bait on warm, clear days. Do not graze animals in treated areas.

**ORCHARDS, VINEYARDS, NURSERIES, NON-BEARING FRUIT TREES, ORNAMENTALS RESTRICTIONS FOR USE:** Use RIDALL-Z to control Meadow Voles (*Microtus pennsylvanicus*) and Pine Voles (*M. pinetorum*) in orchards, vineyards, non-bearing fruit trees, and ornamentals. Bait applications should be made during the non-bearing season when the grass is not yet matted.

**BAITING:** Provide bait at 2-5 locations on trails or near burrow openings adjacent to damaged trees (one teaspoon per bait point). Do not make broadcast application of bait over growing crops when the bait may lodge on the plants. Do not apply the bait to bare ground. Bait at the rate of 4 pounds per acre in vole infested areas. If a trailbuilder is used, apply one teaspoon of bait at 5 foot intervals in the artificially made runway around infested trees. If heavy rains occur after baiting or additional mice move into area, additional treatment may be necessary. For aerial application apply at the rate of 4 lbs per acre. As with broadcasting do not apply on bare ground or growing crops.

**SUGARCANE**

**RESTRICTIONS FOR USE:** Use RIDALL-Z to control Roof Rats (*Rattus rattus*), Rice Rats (*Oryzomys palustris*), and Cotton Rats (*Sigmodon hispidus*) in Florida and Roof Rats, Norway Rats (*R. norvegicus*), and Polynesian Rats (*R. exulans*) in Hawaii.

**BAITING:** Apply untreated rolled oats or barley (Prebait) at the rate of 5 lbs per acre to condition rats to extra food supply 2-4 days before baiting. This will increase acceptance of the bait. RIDALL-Z bait material must not be applied to sugarcane that is less than 6 months of age nor within 30 days of harvest. The total number of bait applications must not exceed four, and the total bait applied shall not exceed 20 lbs per acre. Broadcast the bait with a ground spreader or an aircraft at the rate of 5 lbs per acre.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this product when such use and/or handling is contrary to label instructions.

**RESTRICTED USE  
PESTICIDE**

For retail sale and use only, by Certified Applicators or persons under their direct supervision and only for the uses covered by the Certified Applicators Certification.

**Chempar's**

**RIDALL-Z™**

**Rodent Field & Agricultural Bait**

For use in the Control of:

- Rats in sugarcane
- Ground Squirrels (noncrop use)
- Prairie Dogs (Grasslands)
- Pine & Meadow Voles (Orchards & noncrop areas)
- Gophers and Moles

Active Ingredient	1.88%
Zinc Phosphide	98.12%
Inert Ingredients	
<b>TOTAL</b>	<b>100.00%</b>

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

- Any persons applying zinc phosphide products and experiencing signs and symptoms such as nausea, abdominal pain, lightness in the chest, or weakness, should see a doctor immediately.
- If swallowed, drink 1-2 glasses of water and induce vomiting by touching the back of the throat with the finger. Avoid the use of oils. Have person lie down and keep warm. Do not induce vomiting or give anything by mouth to an unconscious person.
- If on skin or eyes, flush with plenty of water.
- See left side for additional precautionary statement.

Manufactured by:

**Chempar**

**A DIVISION OF  
LIPHA CHEMICALS, INC.**

**880 MADISON AVENUE  
NEW YORK, NY 10021**

**EPA Reg. No. 7173-195**

**EPA EST. NO. 7173-NY-01**

**NET CONTENTS: 50 Lbs. (22.73 Kgs.)**

**Product No. 8000  
Label No. 5088-0885**

**RESTRICTIONS FOR USE (cont)**

**RANGELANDS**

**RESTRICTIONS FOR USE FOR CONTROL OF PRAIRIE DOGS:** Use RIDALL-Z to control Blacktailed Prairie Dogs (*Cynomys ludovicianus*) and Whittailed Prairie Dogs (*C. leucurus*) on ranges in the western U.S. (Colorado, Montana, Arizona, New Mexico, Texas, Oklahoma, Kansas, Nebraska, North & South Dakota). Use this bait only in areas not inhabited by Black Footed Ferrets. Consult Federal, Local, and State Wildlife authorities concerning the range of the Ferret before using RIDALL-Z.

**BAITING:** Use a prebait of one teaspoon of untreated rolled oats or barley placed near each active mound 2-4 days before using RIDALL-Z bait. This will increase prairie dog acceptance of the bait. When 50% of the prebait is eaten, hand bait using one teaspoon of bait placed within a one foot radius of each mound entrance. The recommended baiting time is from July to December with only one application each year. Bury the uneaten bait or dead animals found.

**FOR CONTROL OF GROUND SQUIRRELS:** Use RIDALL-Z bait to control California Ground Squirrels (*Spermophilus beecheyi*) and Richardson's Ground Squirrels (*S. richardsoni*) in the western US.

**BAITING:** Prebait with one teaspoon of rolled oats or barley near each active burrow system. After 2-4 days apply RIDALL-Z at the rate of one teaspoon of bait near each active burrow. Place the bait on bare soil. Apply bait in the mid-afternoon so most squirrels will expire inside the burrows. The recommended baiting time is late spring and early summer (April-June) as soon as squirrels emerge from hibernation.

**NON-CROP AREAS**

**RESTRICTIONS FOR USE-HAND BAITING:** Use RIDALL-Z to bait areas such as right-of-ways, golf courses, nurseries and lawns to control California Ground Squirrels (*Spermophilus beecheyi*), Richardson's Ground Squirrels (*S. richardsoni*), Thirteen-lined Ground Squirrels (*S. tridecemlineatus*), Meadow Voles (*Microtus pennsylvanicus*) and Pine Voles (*M. pinetorum*). Do not apply bait over areas where plants are used as food or feed, on roads or over water (such as lakes).

**BAITING:** For ground squirrel control, use untreated rolled oats or barley and place one teaspoon of grain near active burrows 2-4 days before baiting. Apply one teaspoon of RIDALL-Z near active burrows or runways directly on bare ground. Bury all dead animals found and the uneaten bait. For Meadow and Pine Vole control provide bait in 2-5 locations on trails or near burrow openings adjacent to damaged trees use one teaspoon per bait point. Do not apply to bare ground. Bait at the rate of 4 lbs per acre in infested areas. If a trailbuilder is used, apply RIDALL-Z at the rate of one teaspoon of bait at 5-foot intervals in the artificially made runway around infested trees. If heavy rains occur after baiting or additional mice move into area, additional treatment may be necessary.

**BROADCAST BAITING:** Use RIDALL-Z along non-crop right-of-ways to control California Ground Squirrels, Richardson's

Ground Squirrels, and Thirteen lined Ground Squirrels along field borders or ditch banks during a 3-month period after squirrels emergence from hibernation (April-June). Use rolled oats or barley as a prebait and apply over the area at a rate of 4 lbs per acre 2-4 days before providing RIDALL-Z. This should increase bait acceptance. Broadcast in 20 ft swaths using hand or ground driven equipment at a rate not to exceed 4 lbs per acre. Apply bait in the mid-afternoon so most animals will expire in their burrows. Do not broadcast near homes where food or feed is grown or on roads. Consult wildlife authorities for local regulations. In right-of-ways bait may only be applied between the bottom of the ditch and the property. Always bury excess bait left in the equipment. For control of Meadow and Pine Voles, prebait the infested area by broadcasting rolled oats or barley at the rate of 4 lbs per acre 2-4 days before applying the bait. Using hand or mechanically driven equipment, apply RIDALL-Z in 20-ft swaths at a rate not to exceed 4 lbs per acre.

**GOPHERS AND MOLES**

**RESTRICTIONS FOR USE:** Use RIDALL-Z to control the Plains Pocket Gopher (*Geomys bursarius*), Southeastern Pocket Gopher (*G. pinetis*), Yellow-faced Pocket Gopher (*Pappageomys castanops*), Valley Pocket Gopher (*Thomomys bottae*), and the Northern Pocket Gopher (*T. talpoides*) in non-crop areas such as right-of-ways, golf courses, nurseries and lawns, on rangelands, and in forests in the central and western US. Similarly RIDALL-Z may also be used to control Eastern Moles (*Scalopus aquaticus*), Star-nose Moles (*Condylura cristata*), California Moles (*Scapanus latimanus*), and Townsend Moles (*S. townsendi*) in Eastern, Southern, and Western US. Do not apply this bait in gardens and areas where food or feed may be contaminated. Only apply the bait underground. Collect and bury all dead animals found above the ground.

**BAITING:** Locate fresh activity (push piles, earth plugs, or network of surface ridges in the turf). Use a mechanical or metal probe and push it into the ground to locate the underground runway. Place one teaspoon of bait inside the tunnel or uncover the burrow opening and place the bait directly inside. Cover the opening with a rock or leaf litter. Loose soil. Surface baiting will be of little value. It is best to control gophers and moles in the late summer and early fall or when burrowing activity is fresh. With hand baiting or if a burrow builder is used, apply the bait at a rate not to exceed 4 lbs per acre. Several days after applying bait, open burrows previously baited and leave open. If gophers or moles remain, they should plug the holes closed within one day. If this occurs, then the application should be repeated in the same manner.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store in original container only in a locked storage area inaccessible to children and pets. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

7173-195 PM 16  
171

RECEIVED  
30 1986  
7173-195

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION**

Keep away from humans, domestic animals and pets. Do not breathe fumes or dust. May be fatal if swallowed. Do not allow contact with skin. Wash hands after use. Do not use near feed or foodstuffs. Keep away from children. Wear gloves and protective clothing when using bait. Do not use or store in house.

**ENVIRONMENTAL HAZARDS**

This product is toxic to wildlife and fish. Birds and other wildlife feeding on treated bait may be killed. Do not apply to open ground, roads, and areas in which no vegetation occurs. Pick up and properly dispose of all spilled toxic bait. Keep out of lakes, streams, ponds, and aquatic systems. Do not apply where runoff from this product will enter an aquatic system. Check for and dispose of excess spoiled bait and carcasses daily until the bait is expended or removed.

**ENDANGERED SPECIES  
CONSIDERATIONS**

Before field use of this product, consult Federal, State, and Local game authorities to determine if the following wildlife are found in the area. If so, observe the following:

- Whooping Crane (*Grus americana*)  
Do not use this product 30 days prior to arrival and 30 days after the whooping crane leaves known crane habitat.
- Attwater's Greater Prairie Chicken (*Tympanuchus cupido attwateri*)  
Do not use this product in the presently occupied range of the Attwater greater prairie chicken in the following Texas counties, Arkansas, Austin, Brazoria, Colorado, Dowlit, Fort Bend, Galveston, Galind, Harris, Refugio, Victoria, Walker, and Wharton.
- Yellow-Shouldered Black Bird (*Agelaius xanthomus*) and Puerto Rican Plain Pigeon (*Columba inornata wetmorei*)  
This product must not be used in the following areas except in tamper proof bait boxes (within 10 kilometers of central Aguera, Lago Cidra, Ceiba, San German, 15 Kilometers of La Esperanza, south of Highway 2 from city of Mayaguez to the city of Ponce and all of Mona Island, Puerto Rico).
- Utah Prairie Dog (*Cynomys parvidens*)  
Do not use this product in the range of the Utah Prairie Dog (Utah).
- Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*)  
Do not use this product within 1/2 mile of salt marsh vegetation and/or brackish water wetlands which are located (1) near or adjacent to San Pablo Bay and San Francisco Bay or (2) in the Sacramento River below or adjacent to the confluence of the Sacramento River and the San Joaquin River (California).

**ENDANGERED SPECIES CONSIDERATIONS (cont)**

- Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*)  
Do not use this product within 2 1/2 miles of Baywood Park which is located on Morro Bay (California).
- Aleutian Canada Goose (*Brania canadensis leucopara*)  
Do not use this product from October to March in the Federally closed goose hunting areas in the state of California. See current Duck & Goose Hunting game laws for current map.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This bait must not be applied on roads, near residential areas, or over water. Do not broadcast over growing crops other than sugarcane. Apply bait on warm, clear days. Do not graze animals in treated areas.

**ORCHARDS, VINEYARDS, NURSERIES,  
NON-BEARING FRUIT TREES, ORNAMENTALS**

**RESTRICTIONS FOR USE:** Use RIDALL-Z to control Meadow Voles (*Microtus pennsylvanicus*) and Pine Voles (*M. pinetorum*) in orchards, vineyards, non-bearing fruit trees, and ornamentals. Bait applications should be made during the non-bearing season when the grass is not yet matted.

**BAITING:** Provide bait at 2-5 locations on trails or near burrow openings adjacent to damaged trees (one teaspoon per bait point). Do not make broadcast application of bait over growing crops when the bait may lodge on the plants. Do not apply the bait to bare ground. Bait at the rate of 4 pounds per acre in vole-infested areas. If a trailbuilder is used, apply one teaspoon of bait at 5 foot intervals in the artificially made runway around infested trees. If heavy rains occur after baiting or additional mice move into area, additional treatment may be necessary. For aerial application apply at the rate of 4 lbs per acre. As with broadcasting do not apply on bare ground or growing crops.

**SUGARCANE**

**RESTRICTIONS FOR USE:** Use RIDALL-Z to control Roof Rats (*Rattus rattus*), Rice Rats (*Oryzomys palustris*), and Cotton Rats (*Sigmodon hispidus*) in Florida and Roof Rats, Norway Rats (*R. norvegicus*), and Polynesian Rats (*R. exulans*) in Hawaii.

**BAITING:** Apply untreated rolled oats or barley (Prebait) at the rate of 5 lbs. per acre in cotton rats to extra food supply 2-4 days before baiting. This will increase acceptance of the bait. RIDALL-Z bait material must not be applied to sugarcane that is less than 6 months of age not within 30 days of harvest. The total number of bait applications must not exceed four, and the total bait applied shall not exceed 20 lbs. per acre. Broadcast the bait with a ground spreader or an aircraft at the rate of 5 lbs. per acre.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**RESTRICTED USE  
PESTICIDE**

For retail sale and use only, by Certified Applicators or persons under their direct supervision and only for the uses covered by the Certified Applicators Certification.

**Chempar's  
RIDALL-Z<sup>TM</sup>**

**Rodent Field & Agricultural Bait**

**For use in the Control of:**

- Rats in sugarcane
- Ground Squirrels (noncrop use)
- Prairie Dogs (Grasslands)
- Pine & Meadow Voles (Orchards & noncrop areas)
- Gophers and Moles

Active Ingredient	1.88%
Zinc Phosphide	98.12%
Inert Ingredients	
<b>TOTAL</b>	<b>100.00%</b>

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

- Any persons applying zinc phosphide products and experiencing signs and symptoms such as nausea, abdominal pain, tightness in the chest, or weakness, should see a doctor immediately.
- If swallowed, drink 1-2 glasses of water and induce vomiting by touching the back of the throat with the finger. Avoid the use of oils. Have person lie down and keep warm. Do not induce vomiting or give anything by mouth to an unconscious person.
- If on skin or eyes, flush with plenty of water.
- See left side for additional precautionary statement.

Manufactured by:



**A DIVISION OF  
LIPHA CHEMICALS, INC.  
860 MADISON AVENUE  
NEW YORK, NY 10021**

EPA Reg No. 7173-195  
EPA EST NO. 7173-NY-01

**NET CONTENTS: 50 Lbs. (22.73 Kgs.)**

Product No. 8000  
Label No. 5088-0885

**DIRECTIONS FOR USE (cont)**

**RANGELANDS**

**RESTRICTIONS FOR USE FOR CONTROL OF PRAIRIE DOGS:** Use RIDALL-Z to control Blacktailed Prairie Dogs (*Cynomys ludovicianus*) and Whitetailed Prairie Dogs (*C. leucurus*) on ranges in the western U.S. (Colorado, Montana, Arizona, New Mexico, Texas, Oklahoma, Kansas, Nebraska, North & South Dakota). Use this bait only in areas not inhabited by Black Footed Ferrets. Consult Federal, Local, and State Wildlife authorities concerning the range of the Ferret before using RIDALL-Z.

**BAITING:** Use a prebait of one teaspoon of untreated rolled oats or barley placed near each active mound 2-4 days before using RIDALL-Z bait. This will increase prairie dog acceptance of the bait. When 50% of the prebait is eaten, hand bait using one teaspoon of bait placed within a one foot radius of each mound entrance. The recommended baiting time is from July to December with only one application each year. Bury the uneaten bait or dead animals found.

**FOR CONTROL OF GROUND SQUIRRELS:** Use RIDALL-Z bait to control California Ground Squirrels (*Spermophilus beecheyi*) and Richardson's Ground Squirrels (*S. richardsoni*) in the western U.S.

**BAITING:** Prebait with one teaspoon of rolled oats or barley near each active burrow system. After 2-4 days apply RIDALL-Z at the rate of one teaspoon of bait near each active burrow. Place the bait on bare soil. Apply bait in the mid-afternoon so most squirrels will expire inside the burrows. The recommended baiting time is late spring and early summer (April-June) as soon as squirrels emerge from hibernation.

**NON-CROP AREAS**

**RESTRICTIONS FOR USE-HAND BAITING:** Use RIDALL-Z to bait areas such as right-of-ways, golf courses, nurseries and lawns to control California Ground Squirrels (*Spermophilus beecheyi*), Richardson's Ground Squirrels (*S. richardsoni*), Thirteen-lined Ground Squirrels (*S. tridecemlineatus*), Meadow Voles (*Microtus pennsylvanicus*) and Pine Voles (*M. pinetorum*). Do not apply bait over areas where plants are used as food or feed, on roads or over water (such as lakes).

**BAITING:** For ground squirrel control, use untreated rolled oats or barley and place one teaspoon of grain near active burrows 2-4 days before baiting. Apply one teaspoon of RIDALL-Z near active burrows or runways directly on bare ground. Bury all dead animals found and the uneaten bait. For Meadow and Pine Vole control provide bait in 2-5 locations on trails or near burrow openings adjacent to damaged trees use one teaspoon per bait point. Do not apply to bare ground. Bait at the rate of 4 lbs. per acre in infested areas. If a trailbuilder is used, apply RIDALL-Z at the rate of one teaspoon of bait at 5-foot intervals in the artificially made runway around infested trees. If heavy rains occur after baiting or additional mice move into area, additional treatment may be necessary.

**BROADCAST BAITING:** Use RIDALL-Z along non crop right-of-ways to control California Ground Squirrels, Richardson's

Ground Squirrels, and Thirteen-lined Ground Squirrels along held borders or ditch banks during a 3-month period after squirrels emergence from hibernation (April-June). Use rolled oats or barley as a prebait and apply over the area at a rate of 4 lbs. per acre 2-4 days before providing RIDALL-Z. This should increase bait acceptance. Broadcast in 20 ft. swaths using hand or ground driven equipment at a rate not to exceed 4 lbs. per acre. Apply bait in the mid-afternoon so most animals will expire in their burrows. Do not broadcast near homes where food or feed is grown or on roads. Consult wildlife authorities for local regulations. In right-of-ways bait may only be applied between the bottom of the ditch and the property. Always bury excess bait left in the equipment. For control of Meadow and Pine Voles, prebait the infested area by broadcasting rolled oats or barley at the rate of 4 lbs. per acre 2-4 days before applying the bait. Using hand or mechanically driven equipment, apply RIDALL-Z in 20-ft swaths at a rate not to exceed 4 lbs. per acre.

**GOPHERS AND MOLES**

**RESTRICTIONS FOR USE:** Use RIDALL-Z to control the Plains Pocket Gopher (*Geomys bursarius*), Southeastern Pocket Gopher (*G. pinalis*), Yellow-faced Pocket Gopher (*Pappageomys castanops*), Valley Pocket Gopher (*Thomomys bottae*), and the Northern Pocket Gopher (*T. talpoides*), in non-crop areas such as right-of-ways, golf courses, nurseries and lawns, on rangelands and in forests in the central and western U.S. Similarly RIDALL-Z may also be used to control Eastern Moles (*Scalopus aquaticus*), Star-nosed Moles (*Condylura cristata*), California Moles (*Scapanus latimanus*), and Townsend Moles (*S. townsendi*) in Eastern Southern, and Western U.S. Do not apply the bait in gardens and areas where food or feed may be contaminated. Only apply the bait underground. Collect and bury all dead animals found above the ground.

**BAITING:** Locate fresh activity (push piles earth plugs, or network of surface ridges in the turf). Use a mechanical or metal probe and push it into the ground to locate the underground runway. Place one teaspoon of bait inside the tunnel or uncover the burrow opening and place the bait directly inside. Cover the opening with a rock or leaf. Use loose soil. Surface baiting will be of little value. It is best to control gophers and moles in the late summer and early fall or when burrowing activity is fresh. With hand baiting or if a burrow builder is used, apply the bait at a rate not to exceed 4 lbs. per acre. Several days after applying bait, open burrows previously baited and leave open. If gophers or moles remain, they should plug the holes closed within one day. If this occurs, then the application should be repeated in the same manner.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by Storage or Disposal.  
**STORAGE:** Store in original container only in a locked storage area inaccessible to children and pets.  
**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.  
**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

09.093

00.013

ACCEPTED  
MAY 30 1986  
EPA Reg. No. 7123-195