

7173-215

5/30/2001

1/1

**PRECAUTIONARY STATEMENTS**  
**Hazard to Humans and Domestic Animals**

**CAUTION:** Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding. Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking.

**FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Note to Physician or Veterinarian: If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**STORAGE AND DISPOSAL:**

Do not contaminate water, food or feed by storage or disposal. Store in original container in a locked storage area.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Dispose of empty container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 7173-215  
 EPA Est. No. 7173MM-1

NET WEIGHT: 1 Pound (454 g)

**ACCEPTED**  
 MAY 30 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7173-215

# MAKI

## MINI BLOCKS

### With Bitrex

**KILLS NORWAY RATS,  
 ROOF RATS, HOUSE MICE AND  
 WARFARIN-RESISTANT NORWAY RATS**

**NORWAY RATS MAY CONSUME A  
 LETHAL DOSE IN ONE FEEDING  
 WITH FIRST DEAD RODENTS  
 APPEARING FOUR OR FIVE DAYS  
 AFTER TREATMENT BEGINS**

Active Ingredient: Bromadiolone .....	0.005%
Inert Ingredients .....	99.995%
Total .....	100.000%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking. **If swallowed, call a physician at once. See side panel for additional precautionary statements.**

**LIPHATECH**

Manufactured By:  
 LIPHATECH, INC.  
 3101 W. Custer Avenue  
 Milwaukee, WI 53209

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** For control of Norway Rats, Roof Rats and House Mice.

**Urban areas:** May be used in and around the periphery of homes, industrial, commercial and public buildings. This product may be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. May also be used in alleys. Do not use in sewers.

**Non-urban areas:** May be used inside homes and agricultural buildings.

Do not place bait where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors or walls, or locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:**

**Rats:** Use 3 to 16 mini blocks per placement. Space placements at intervals of 15 to 30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for 10 days or until fresh signs of rat activity no longer are evident.

**Mice:** Use one mini block per placement. At points of very high mouse activity, two mini blocks may be used. Space placements at intervals of 8 to 12 feet. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity no longer are evident.

**Rats/Mice:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.