



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

7173-309

Date of Issuance:

11/1/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Rozol® Field Rodent Bait

Name and Address of Registrant (include ZIP Code):

Kylli Paavola
Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).


Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Sincerely, 
Gene Benbow, Product Manager 7
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

Date:

11/1/22

EPA Form 8570-6

2. You are required to comply with the data requirements described in the generic data call-in (GDCI) identified below:

a. Chlorophacinone GDCI-067707-1616

You must comply with all of the data requirements within the established deadlines. If you have questions about the GDCI listed above, you may contact the Chemical Review Manager in the Pesticide Re-Evaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 7173-309.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 10/01/2020

If you have any questions, please contact Ralph Narain at 202-566-2853 or at Narain.Ralph@epa.gov.

Enclosure


RESTRICTED USE PESTICIDE
DUE TO HAZARD TO NONTARGET ORGANISMS
For retail sale to and use by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.

Rozol® Field Rodent Bait

(Alternate Brand Name: Rozol® FR)

FOR CONTROL OF POCKET GOPHERS (*Thomomys* spp. and *Geomys* spp.), CALIFORNIA GROUND SQUIRRELS (*Spermophilus beecheyi*), RICHARDSON'S GROUND SQUIRRELS* (*Uroditellus richardsonii*), AND COLUMBIAN GROUND SQUIRRELS* (*U. columbianus*)

Active Ingredient: chlorophacinone 0.005%
Other Ingredients 99.995%
Total 100.000%



KEEP OUT OF REACH OF CHILDREN

CAUTION:

See side [back] [inner] [outer] panel [carton] [label] for additional precautionary statements.

(Liphatech Logo)
Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(800) 351-1476

Net Weight: 4 pounds (lbs) up to 2000 pounds (lbs)
This product cannot be sold or distributed in packaging that holds less than 4 pounds of bait

*Use for control of these species not permitted in California

EPA Est. No. 7173-WI-1
EPA Reg. No. 7173- GNO

Product No.:
Label No.:

FIRST AID:	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin with plenty of cool water for 15-20 minutes. • Call a poison control center or doctor immediately for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor immediately for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor immediately for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
TREATMENT FOR PET POISONING: If animal eats bait, call veterinarian at once.	
<p>NOTE TO PHYSICIAN OR VETERINARIAN: Have this label with you when obtaining treatment advice. Contains the anticoagulant chlorophacinone. If swallowed, this material may reduce the clotting ability of blood and cause bleeding. For humans or animals ingesting bait and/or showing obvious poisoning symptoms (bleeding or prolonged prothrombin times), give Vitamin K₁ intramuscularly or orally.</p>	

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

CAUTION. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Keep away from children, domestic animals and pets.

Personal Protective Equipment (PPE) Requirements

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.
- Waterproof gloves.

User Safety Recommendations

Users should:

- Wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Wash the outside of gloves before removing them.
- Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to mammals, birds, fish, and freshwater invertebrates. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: It is against Federal law to use any pesticide in a manner that results in the death of an endangered species. Use of this product may pose a hazard to endangered or threatened species. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. To obtain Bulletins, no more than six months before using the product, consult <http://www.epa.gov/espp/> or call 844-447-3813. You must use the Bulletin valid for the month in which you will apply the product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL and follow all use directions and use precautions. Only use this product for the pests, sites, and application methods specified on this label.

POCKET GOPHERS


USE RESTRICTIONS: Use only to control pocket gophers (*Thomomys spp.* and *Geomys spp.*) in lawns, golf courses, forest areas, tree and forestry plantations, nurseries, fallow agricultural land, non-crop areas, fence rows, highway medians and rights-of-way, rail and utility rights-of-way, ditches, canal banks and levees, ornamental gardens, parks, campgrounds and other recreational areas. Bait must be applied directly into pocket gophers' burrow systems. Only apply bait underground.

Application Directions: Burrowing pocket gophers throw out low, fan-shaped mounds on either side of the entrance to their underground tunnel. The entrance holes are on the flat side of the fan and are plugged with loose soil.

Treatment: Bait placement can be made in one or both of the following ways:

1. Remove the plug on the flat side of the fan with a shovel or long-handled tablespoon. Ensure plug is removed all the way to the main run which runs parallel to the ground surface. Place 1/2 cup of bait as far down into the hole as possible. Re-close the opening, using care not to cover the bait with soil.
2. Using a metal rod, probe 6–12 inches deep to locate the main tunnel. Consult the diagram below for location to probe. Drop 1/2 cup of bait into the tunnel and cover the hole so light cannot enter the tunnel system.

Make 2–3 bait placements per burrow system. Maintain a constant supply of bait in the burrow system for as long as there is gopher activity. Do not apply bait on the soil surface.



The right and wrong ways to use a probe for placing pocket gopher bait are shown. Bait should be placed loosely in the main runway - not in the lateral tunnels or imbedded in the bottom of the runway. For additional information, consult the Federal and State rodent control bulletins for a full discussion of pocket gopher burrowing habits.

Machine application using burrow builder equipment: Follow manufacturer's instructions for the equipment used to apply this bait. Make artificial burrows at the same depth as natural burrows about 15 to 30 feet apart. Apply 1.5 to 8 pounds of bait per acre. Use the high rate in locations with greater pocket gopher activity. Apply when soil conditions allow formation of good artificial burrows. Do not treat the entire field. Apply as:

1. A targeted treatment to the part of the field where pocket gophers have established burrows. Make 6 to 8 rows in the area infested with pocket gophers. Do not exceed 4 applications per year.
2. Perimeter treatment to reduce potential migration of pocket gophers into the field. Make 3 rows around the perimeter of the field. Use only in locations where there is a risk of pocket gopher damage. Do not exceed 3 applications per year. For best results apply in the fall, spring, and mid-summer.

GROUND SQUIRRELS

USE RESTRICTIONS: This bait may be used only for control of California ground squirrels (*Spermophilus beecheyi*), Richardson's ground squirrels* (*Uroditellus richardsonii*), and Columbian ground squirrels* (*U. columbianus*) in lawns, golf courses, vineyards, pome fruit, stone fruit, citrus fruit, avocado and nut tree orchards, berry and small fruit, tropical and subtropical fruit (edible and inedible peel), herbs and spices, forest areas, tree and forestry plantations, nurseries; grass forage, fodder and hay; nongrass animal feeds (forage, fodder, straw and hay); forage, fodder, and straw of cereal grains; cereal grains; edible seed crops, oil seed crops, fiber crops, vegetable crops, pastures and rangeland, fallow agricultural land, non-crop areas, fence rows, highway medians and rights-of-way, rail and utility rights-of-way, ditches, canal banks and levees, ornamental gardens, parks, campgrounds and other recreational areas. Applications within the above sites may be made year-round if using tamper-resistant bait stations See each application method below for allowed use sites and restrictions.

Do not apply within 50 feet of any body of surface water. Do not apply this product in or within 50 feet of homes or other human residences. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treatment area during application.

Do not apply on roads, in water or to walkways. When applying bait along canals or other waterways, use equipment that directs bait away from water. Do not apply directly to food or feed crops, application may be made along fence lines, border areas and buffer strips adjacent to crops. Do not apply bait by aircraft.

*Use for control of these species not permitted in California: Richardson's ground squirrel and Columbian ground squirrel.

Application Directions: Ground squirrels prefer to eat fresh green grass and other plants. When these are not available (early spring or when green vegetation dries up in the hot summer), the ground squirrel's diet converts to grain-based foods and they will readily consume this grain-based bait. Before applying, test for the ground squirrels' acceptance of grain-based foods. Scatter some raw oats or wheat on a portion of the infested land and mark the test area with a flag or other identification. Observe the area to confirm only ground squirrels are consuming the raw oats or wheat. Do not apply the bait if it appears that nontarget species are consuming the test grain. For best results apply bait by broadcast or spot baiting when fresh green ground cover is not present on the majority of the treatment area. Monitor the area to ensure that nontarget species are not consuming the bait or ground squirrels. For best results apply in early spring immediately after ground squirrels emerge from winter hibernation.

Bait station baiting (For California, Richardson's, and Columbian ground squirrels): Bait station applications may be made in lawns, golf courses, vineyards, pome fruit, stone fruit, citrus fruit, avocado and nut tree orchards, berry and small fruit, tropical and subtropical fruit (edible and inedible peel), herbs and spices, forest areas, tree and forestry plantations, nurseries; grass forage, fodder and hay; nongrass animal feeds (forage, fodder, straw and hay); forage, fodder, and straw of cereal grains; cereal grains; edible seed crops, oil seed crops, fiber crops, vegetable crops, pastures and rangeland, fallow agricultural land, non-crop areas, fence rows, highway medians and rights-of-way, rail and utility rights-of-way, ditches, canal banks and levees, ornamental gardens, parks, campgrounds and other recreational areas.

Apply bait in secured tamper-resistant bait stations. Fill bait stations with 1 to 4 pounds of bait, and secure the stations near active burrows, runways, and other locations where signs of ground squirrel activity are observed. Space stations 20 to 100 feet apart, depending on the density of the ground squirrel population. Check stations at least weekly to ensure an uninterrupted supply of fresh bait for at least 30 days or until all signs of ground squirrel feeding have stopped. When feeding by ground squirrels ceases, and there is no longer a threat of continued reinvasion, remove bait from bait stations and dispose of properly. Bait stations may be kept in place permanently but should be loaded with toxic bait only when ground squirrels are readily accepting grain-based foods. Do not exceed 4 pounds of bait per station.

Spot baiting (For California and Richardson's ground squirrels only): Spot baiting applications may only be made in forest areas, tree and forestry plantations, nurseries, fallow agricultural land, non-crop areas, fence rows, highway medians and rights-of-way, rail and utility rights-of-way, ditches, canal banks and levees.

Scatter ¼ cup of bait (approximately 2 ounces) over 40 to 50 square feet (a circle approximately 7 to 8 feet diameter) near burrows and active runways. Do not pile the bait. Make a second application 3 to 4 days after the first, following the same procedure as the first application. If ground squirrel activity persists, a third application may be made 3 to 4 days later. An uninterrupted supply of bait should be available for 6 to 8 days.

Do not exceed 10 pounds of bait per acre per application. Do not exceed 20 pounds of bait per acre per calendar year. Do not graze livestock or plant food or feed crops in spot-treated areas while bait is present. Do not harvest agricultural crops from treated fallow agricultural land for 360 days after treatment. In the state of California, spot baiting placements must be covered with grass or a shingle to avoid exposing to nontarget species.

Broadcast baiting (For California and Richardson's ground squirrels only): Broadcast baiting may only be made in forest areas, tree and forestry plantations, nurseries, fallow agricultural land, non-crop areas, fence rows, highway medians and rights-of-way, rail and utility rights-of-way, ditches, canal banks and levees.

Using a commercial spreader, apply up to 10 pounds of bait per swath acre, in the area of ground squirrel activity. Make a second application 4 days after the first. If necessary, make a third application four days after the second application, in order to maintain an uninterrupted supply of bait for 6 to 8 days.

Do not exceed 10 pounds of bait per acre per application. Do not exceed 20 pounds of bait per acre per calendar year. Do not graze livestock or plant food or feed crops in broadcast areas while bait is present. Do not harvest agricultural crops from treated fallow agricultural land for 360 days after treatment. Do not broadcast bait on rangeland. Do not apply by broadcast in the state of California.

Clean-up: Clean up any spilled bait or bait in piles for reuse or disposal according to label "Pesticide Disposal" instructions. Use detergent and hot water to wash spoons, cups and other application equipment. Do not use the same equipment for mixing, holding or transferring food or feed.

Surveillance and Follow-Up: Dead pocket gophers and ground squirrels should begin to appear in or near treated areas within 4 to 5 days after the first bait application. Visit the application site to monitor the effects of treatment and to collect and dispose of dead rodents and search for nontarget animals 4 days after the first application and at subsequent intervals of 1 to 2 days for at least 2 weeks after the last bait application, or longer if carcasses are still being found. Collect and dispose of carcasses in the late afternoon, near sundown, to reduce the potential for nocturnal scavengers or predators to find carcasses on the ground surface. Wearing gloves, collect and properly dispose of visible carcasses of pocket gophers, ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Wear gloves and use leakproof plastic bags or other suitable containers for transporting carcasses not buried on site.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of in trash or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. **[Plastic:]** Completely empty container, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill. **[Paper:]** Completely empty container, then dispose of empty container in trash or at an approved waste disposal facility.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of product when such use and/or handling is contrary to label instructions.

Optional Text/ Pictures/ Graphics:

Moisture Resistant

Effective control

Palatable bait

Satisfaction guaranteed or your money back.

Manufactured in the USA with globally and domestically sourced materials.

Optional Text/ Pictures/ Graphics Continued:

[California Ground Squirrel]



[Richardson's Ground Squirrel]



[Columbian Ground Squirrel]




[Pocket Gopher]




[3 x 3 Perimeter Baiting Diagram for Burrow Builder Pocket Gopher Application]



[Mechanical Burrow Builder Schematic]



[Mechanical Burrow Baiting Diagram – Perpendicular Burrows]



[Mechanical Burrow Builder]

