

Net Contents: 5 Gallons

EPA EST.

Sustar 2-S

PLANT GROWTH REGULATOR

MADE IN U.S.A.



CAUTION

KEEP OUT OF REACH OF CHILDREN

Active Ingredient	30%
Diethanolamine salt of N-(4-methyl-3-[(1,1,1-trifluoroethyl)sulfonyl]amino)phenyl)acetamide†	
Inert Ingredients	70%
*Equivalent to 2.0 pounds (22% by weight) of N-(4-methyl-3-[(1,1,1-trifluoroethyl)sulfonyl]amino)phenyl)acetamide per gallon.	
† U.S. Patent No. 3,639,474	

EPA Reg. No. 7182-3

APPLICATION: SUSTAR 2 S Plant Growth Regulator is useful in noncrop turf areas for retarding grass growth and suppressing seedhead formation of the following species:

Annual Bluegrass	Fescue
Bahiagrass	Kentucky Bluegrass
Bentgrass	Orchardgrass
Bermudagrass	Perennial Ryegrass
Bromegrass	Quaragrass
Dallisgrass	St. Augustinegrass

SUSTAR 2 S is also useful to suppress growth of clovers and sandhorns.

SUSTAR 2 S is recommended to reduce mowing requirements for three to eight weeks and to improve appearance of ornamental turf.

Highway rights of way — especially where regular mowing and maintenance is difficult, costly, or hazardous due to traffic or slope.

Golf courses — on roughs, fairways, and around ditches, at tees, etc., where mowing is difficult or impossible.

Industrial sites — hard to mow areas adjacent to buildings, etc., are heavily obstructed with trees, hydrants, piping, or similar objects; on hillside areas and around parking lots and safety islands.

Cemeteries — around markers and on roadsides.

DIRECTIONS FOR USE: Apply SUSTAR 2 S as a single seasonal treatment shortly before or during active plant growth. SUSTAR 2 S may be applied without mowing, but for a more even appearance, mowing prior to application is recommended. Mix 1 1/2 gallons of SUSTAR 2 S formulation with 40-80 gallons of water and spray evenly on one acre of turf with calibrated ground spray equipment.

IMPORTANT: For best results, SUSTAR 2 S must be applied uniformly. Do not apply to areas covered with grass clippings or leaves. To avoid turf injury, SUSTAR 2 S should not be applied to golf greens, to newly seeded areas, or to newly mowed areas cut to less than one inch in height.

Occasionally SUSTAR 2 S may cause slight temporary discoloration of the turf.

Avoid spraying herbaceous or woody ornamental plants since growth, flowering, and fruit set may be retarded.

After use, rinse spray equipment thoroughly with water.

Prolonged contact with SUSTAR 2 S may accelerate corrosion of metal sprayer parts.

CAUTION: Avoid contact with skin or eyes. In case of contact, immediately flush skin or eyes thoroughly with water. Do not store near food or feed. Do not use on home lawns or forage lands. Do not allow animals to graze on treated areas. Do not reuse empty containers. Rinse with water and discard in a safe place.

NOTICE: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of this manufacturer, no warranty, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with labeled directions. The buyer assumes all responsibility for injury, loss or damage, direct or consequential, resulting from misuse as such, or in combination with other materials.

ACCEPTED
 Feb. 6, 1975
 UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT
 3M Agrichemicals 3M COMPANY
 7182-3

SUSTAR is a registered trademark of 3M Company.