



CAUTION
KEEP OUT OF REACH
OF CHILDREN

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. In case of contact, flush with plenty of water; for eyes, get medical attention. Wear rubber gloves and respirator when mixing Seed Treater with potato seed pieces. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Valley Gro
Potato Seed Treater

E.P.A. REG. NO. 7192-4 **FUNGICIDE** Net Weight 50 Lbs.

ACTIVE INGREDIENT:

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis (dithiocarbamate)] zinc with 1 part by weight (16.1%) ethylenebis [dithiocarbamic acid] bimolecular and trimolecular cyclic anhydrosulfides and disulfides 7.0%

INERT INGREDIENTS 93.0%
100.0%

Zinc expressed as metallic 1.32%
U. S. Patent Pending

Manufactured by SIMPLOT SOIL BUILDERS, East Grand Forks; Minnesota

POTATO-CUT SEED: *Fusarium* Seed Piece Decay, Black Leg, Seed-borne Common Scab - Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed pieces. For effective control, Seed Treater must be applied to entire surface of each piece immediately after cutting. If planting is delayed, follow the recommended storage practices of cut seed pieces in your area.

POTATO-WHOLE SEED: Seed-borne Common Scab - Apply 1 to 1½ pounds of Seed Treater to each 100 pounds of seed. Seed Treater should be applied evenly to entire surface. If the treated whole seed is cut, a second application will be necessary if Black Leg and *Fusarium* Seed Piece Decay are to be controlled.

CAUTION: Do not use treated seed for food or feed purposes. This product is toxic to fish. Do Not contaminate water by cleaning of equipment, or disposal of wastes, and apply this product only as specified on this label.

