

**DIRECTIONS FOR USE**

**POTATOES:** Early blight, late blight - use 2 pounds in 100 gallons of water per acre by ground application or 2 pounds per acre in 3 to 6 gallons of water per acre in aircraft application. Start applications when plants are small (2 to 6 inches). Apply at 5 to 10 day intervals or oftener if weather conditions are conducive for rapid blight development. Blite out may be applied up to date of harvest.

**SUGAR BEETS:** Cercospora Leaf Spot - Use 3 pounds per acre in 100 to 150 gallons of water. Apply when disease first appears in area (usually late July or early August). Repeat at 5 to 10 day intervals as required to maintain control. Do not feed treated tops to meat or dairy animals. Blite out cannot be applied within 10 days of harvest.

# SIMPLOT SOILBUILDERS BLITE OUT

NET WT 50#

- SUGAR BEET & POTATO FUNGICIDE -

**ACTIVE INGREDIENTS:**

EPA Reg. 7192-18

A mixture of 5.2 parts by weight (83.9%) of ammoniates of (ethylenebis (dithiocarbamate)) - zinc with 1 part by weight (16.1%) ethylenebis - (dithiocarbamic acid) bi-molecular and trimolecular cyclic anhydrosulfides and disulfides . . . . . 48.0%  
 Manganese Ethylene Bisdithio-carbamate (Maneb) . . . . . 32.0%  
 INERT INGREDIENTS . . . . . 20.0%  
 100.0%

Zinc expressed as metallic - 8.4%

Manganese expressed as metallic - 8.0%


**WARNING: KEEP OUT OF REACH OF CHILDREN:** May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. In case of contact, flush with plenty of water; for eyes, get medical attention.

Rinse equipment and **DISPOSE OF WASTES** by burying in non-croplands away from water supplies. Container should be disposed of by burying with wastes.

**MANUFACTURED BY:** Simplot Soilbuilders, Box 590,  
East Grand Forks, Minnesota

**CAUTION**

Do not allow this material to drift into neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions, because animal, plant or crop injury, illegal residues or other undesirable results may occur.



**FOLLOW DIRECTIONS CAREFULLY**

Method and timing of application, crop and weather conditions, mixtures with other chemicals not specifically recommended, and other factors are beyond control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the published directions.