

10 1/2 13 1/2 15 1/2 17 1/2 19 1/2 21 1/2 23 1/2 25 1/2 27 1/2 29 1/2 31 1/2 33 1/2 35 1/2 37 1/2 39 1/2 41 1/2 43 1/2 45 1/2 47 1/2 49 1/2 51 1/2 53 1/2 55 1/2 57 1/2 59 1/2 61 1/2 63 1/2 65 1/2 67 1/2 69 1/2 71 1/2 73 1/2 75 1/2 77 1/2 79 1/2 81 1/2 83 1/2 85 1/2 87 1/2 89 1/2 91 1/2 93 1/2 95 1/2 97 1/2 99 1/2 100

**DANGER—POISON—PRECAUTIONS** ☠



**POISONOUS IF SWALLOWED**  
If swallowed, induce vomiting by inducing a gag reflex or by giving warm salt water (1 tablespoon of salt to glass of warm water). Repeat until mouth feels clean. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

**POISONOUS BY SKIN CONTACT**  
If on skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If eye material gets into the eye, wash immediately with running water for at least ten minutes.

**POISONOUS IF BREATHED**  
If warning symptoms appear, take or administer two 2/150-grain atropine tablets immediately. Get medical attention at once. ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below). Have patient lie down and keep quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

**POISONOUS TO FISH AND WILDLIFE**  
This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

**SOUTHERN FARMERS ASSOCIATION**

**DANGER ☠ POISON**  
**ANTIDOTE ATROPINE**

ATROPINE IS THE SPECIFIC ANTIDOTE FOR Methyl Parathion. Give 2 to 4 tablets (2 mg. each) immediately. Repeat every 15 to 30 minutes until the patient is breathing and has a pulse. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

**INTERNAL:** If swallowed, call a physician immediately. Induce vomiting by inducing a gag reflex or by giving warm salt water (1 tablespoon of salt to glass of warm water). Repeat until mouth feels clean. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

**EXTERNAL:** If the material has been applied on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If eye material gets into the eye, wash immediately with running water for at least ten minutes.

**IF WARNING SYMPTOMS APPEAR:** Take or administer two 2/150-grain atropine tablets immediately. Get medical attention at once. ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below). Have patient lie down and keep quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

**NOTE TO PHYSICIAN**

**WARNING SYMPTOMS:** Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

**TREATMENT:** Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine should be injected at 10 minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Exposure to cholinesterase inhibitors may, at times, cause prolonged susceptibility to various doses of any cholinesterase inhibitor. Allow no further exposure until the time for cholinesterase reactivation has been allowed as determined by blood test.

**DANGER**

Do not swallow or breathe dust or fumes. Do not breathe dust or fumes. Wear clean natural rubber gloves, clean, waterproof protective clothing, and goggles. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or drinking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water, including before re-use. In case of spillage, wash with large amounts of water. Decontaminate area and equipment by washing with a concentrated solution of detergent and rinse with water.

**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY BY CERTIFIED APPLICATOR'S CERTIFICATE

**ACCEPTED**  
APR 17 1978  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered by the Administrator under No. 7838-2



**4 LB. Methyl Parathion**  
**Emulsifiable Insecticide Concentrate**

**STOP - READ LABEL BEFORE USE**

**POISON**  
See Side Panels for Antidote & Precautions



**DA**  
Keep Children Off

1 gallon of SFA BRAND 4 LB. METHYL PARATHION contains 4 pounds of Methyl Parathion

|   |       |
|---|-------|
| ACTIVE INGREDIENTS (by weight)              | /     |
| *0,0-dimethyl 0-p-nitrophenyl thiophosphate |       |
| Aromatic Petroleum Solvent (Xylene)         |       |
| INERT INGREDIENTS                           |       |
| Inert (by weight)                           | ..... |
| TOTAL                                       | ..... |
| *Methyl Parathion                           | ..... |

Distributed By:  
**SOUTHERN FARMERS ASSOCIATION**  
824 North Palm Street  
NORTH LITTLE ROCK, ARKANSAS 72119

**NET CONTENTS**

U.