

FOR MAIL

1. NAME _____

2. ADDRESS _____

3. CITY _____

4. STATE _____

5. ZIP _____

6. PHONE _____

7. DATE _____

8. TIME _____

9. FORMAL 57 _____

[illegible]

Bulk Spray Treatment-Peanuts

Apply 2½ pts. of **FOR-MAL 57** in 5 ga
15 tons of farmers' stockpeanuts as the
storage. Apply coarse spray uniformly. Preferably
a suitable mechanical spray applicator which regulates
rate of application to flow of peanuts. Otherwise,
adjust pressure and size of nozzle opening to rate of
flow of peanuts.

After warehouse has been filled, level surface of peanuts before spraying. Apply surface spray using 1 ½ lbs. of premium grade malathion 25% WP in 2 gals. of water and apply at the rate of 2 gals./1000 sq.ft. of surface. Apply first treatment as soon as bin is filled and leveled, but not later than the first week in October. Apply the second surface treatment 1 month later, followed by treatments of 2 month intervals. Use a piston type power sprayer equipped with agitator and nozzle capable of delivering a coarse spray. Repeat surface application 60 days later if warm temperatures prevail. Otherwise delay further treatment until later winter.

FOR-MAL 57

CONTAINS 5 LBS. PREMIUM GRADE MALATHION PER GALLON

CONTENTS

ACCEPTED
JUL 10 1975
Under the Federal Insecticide
Fungicide and Fungicide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 100-100000

SPECIMEN LABEL

CAUTION: Keep out of reach of children.

Source: *Franklin D. Roosevelt Library, Hyde Park, New York*

ANOTHER FINE PRODUCT OF

FORSHAW CHEMICALS, INC.

F

1150 S. A. E. STREET, CHARLOTTE, N. C. 28208

For the first time, the authors have shown that the *in vitro* and *in vivo* results are in good agreement. The *in vitro* results are in good agreement with the *in vivo* results, indicating that the *in vitro* model is a good representation of the *in vivo* situation. The authors have also shown that the *in vitro* model is a good representation of the *in vivo* situation. The authors have also shown that the *in vitro* model is a good representation of the *in vivo* situation.

CAUTION Do not use the following information to determine the amount of product to use. The amount of product to use is determined by the amount of water in the system. The amount of water in the system is determined by the amount of water in the tank. The amount of water in the tank is determined by the amount of water in the tank. The amount of water in the tank is determined by the amount of water in the tank.

CAUTION

[illegible]

empty container. Destroy by perforating
ing. Bury or discard in a safe place away from
supplies.

ANTIDOTE Atropine is antidotal. This is a cholinesterase inhibitor. If swallowed, induce vomiting by giving a tablespoon of salt in a glass of water and repeat until vomit fluid is clear. If spilled on person, remove contaminated clothing and wash skin and arms with soap and water. If spilled in eyes, flush eyes with plenty of water for 10 minutes and get medical attention. Wash clothing before re-use. Call a physician in case of poison symptoms which include nausea, vomiting, salivation, sweating, abdominal cramps, pinpoint non-reactive pupils.

**DO NOT USE, POUR, SPILL, OR STORE NEAR
HEAT OR OPEN FLAME.**

NOTICE-Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

FOR-MAL 57

ACCEPTED
JUN 1965
LIBRARY