CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin. Wash thoroughly after using. Do not get in eyes, on skin or on clothing. For eyes, get medical attention. Avoid contamination of feed and foodstuffs.

ANTIDOTE: Atropine is antidotal. This is a cholinesterase inhibitor.

IF SWALLOWED: Induce vomiting by giving a tablespoon of salt in a glass of water and repeat until vomit is

IF ON SKIN OR CLOTHING: Remove contaminated clothing and wash skin and arms with soap and water.

IF IN EYES: Flush eyes with plenty of water for 10 minutes and get medical attention.

Wash clothing before re-use. Call a physician in case of polson symptons which include nausea, vomiting, salivation, sweating, abdominal cramps, pinpoint nonreactive pupils.

ENVIRONMENTAL HAZARD

Do not contaminate any body of water by application, drift, cleaning of equipment, or disposal of wastes or containers.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Auricultural Extension Service.

Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

FLAMMABLE Keep away from heat and open flame.


DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FORMAL-50 is especially formulated for the preparation of stable emulsions upon mixing with water. These emulsions are easily applied by means of hand power aprayers and by mist blowers. This emulsion is effective in fly control and against many other insects as listed on this label.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Pest-side, spray mixture, or rinstae that cannot be used or chemically processed should be disposed of in a landfill approved for posticides or buried in a safe place away from water supplies. Container Disposal: (a) Reseal container and offer for reconditioning, or (b) Triple rinse (or equivalent) and: offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place. Consult Federal, State or Local Disposal Authorities for ap-



ACTIVE INGREDIENTS:

Malathion (O. O-Dimethyl dithiophosphate of diethyl mercaptosuccinate) 50.00 40.37 Xylene

INERT INGREDIENTS:

9.63

TOTAL 100%

ACCEPTED 110 V 2 : 1978 the see the first of the extension rank do end to the Act. as amended to the persons 7234-36

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Cautions

ANOTHER FINE PRODUCT OF

FORSHAW CHENICALS, INC.

NOTES: Injury may occur on certain ferns including Boston, Malden hair and Pterls, as well as some species of Crassula.

NOTICE

Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ONLY—Mix 2 oz. with one gallon of water. Apply with equipment that can deliver a pin stream of a small amount of FOR-MAL 50 directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housing, junction boxes

APPLICATIONS WITHIN FOOD AREAS OF

FOOD HANDLING ESTABLISHMENTS-

Limited to Crack and Crevice Treatment

or switch boxes, conduits or hollow equipment legs where roaches and ants hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces. Applications of this product in the food areas of food handling , establishments, other than as a crack and crevice treat-

ment, are not permitted.

FLY CONTROL: As a spray, mix FOR-MAL 50 at the rate of 1 gal. per 50 gals, water (1 pt/6 gals). Apply the spray in and around farm buildings and yards and around the exterior of homes and dairy barns thoroughly wetting surfaces such as walls, ceilings, fences and manure piles where files rest such as the shady outside walls of dairy barns; fences and inside and outside of othr farm buildings and structures. NOTE: Remove animals and poultry before applying inside buildings. Do not use where milk is processed and stored. Do not apply on animals or inside dairy barns. Do not use in dwellings. Avoid contamination of feed and feed products.

POTATOES: For the control of aphids on potatoes, make one or more applications as needed of a spray containing 1 pint FOR-MAL 50 with sufficient water to cover one acre. Make no application later than 10 days prior to harvest.

TOMATOES: For the control of spider mites make one or more applications of a spray containing 1-1/2 pts. FOR-MAL 50 with sufficient water to cover one acre. Make no application on tomatoes later than 10 days befure harvest.

PEAS: For control of pea aphid, make one or more applications of a spray containing 1-1/2 pts per acre with no application later than 14 days before harvest.

BEANS: For control of Mexican Bean Beatle and Spider mites, make two or more applications of a spray containing 1-35 pts. FOR-MAL 50 with sufficient water to cover one acre. Make no applications on beans later than 10 days before harvest.

ORNAMENTALS: For control of aphids, spider mites, whitefly, mealybup, thrips, Japanese beetle adult. four-lined leaf bug tarnish plant bug, rose leaf hopper, potato leaf hopper, and back scale crawler on ornamentals. Apply 1-1/2 pts. per 100 gallons water. For control of softbrown scale and Monterey pine scale on ornamentain, apply 2-1/2 pts. FOR-MAL 50 per 100 gais.

Under the Februal Insertion Act.
Fonglicide and Harantist
as amended, for the pesticular
to a state of the pesticular of the pesti