

injury to or loss of desirable trees or other plants may result from failure to observe the following. Do not apply or create equipment near desirable trees or other plants, or in areas where their roots may be sitting, or in locations where the chemical may be washed or moved into contact with them roots. DO NOT use on lawns, walks, driveways, tennis courts or similar areas. Prevent contact with desirable plants. Keep from contact with herbicides insecticides fungicides and acids.

GENERAL INFORMATION

Symptoms of nutrient deficiency generally appear within 3 to 4 weeks for herbaceous plants and 4 to 8 weeks for woody plants after sufficient rainfall has carried the nutrients into the root zone. Chlorosis and subsequent necrosis may occur but succulents plants are less so.

The degree of control and duration of effect will vary with the amount of product applied, soil texture, rainfall, timing of application and other conditions.

Forshaw Chemicals, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risk inherently associated with use of this product. Infection, irritation or other unintended consequences may result because of such factors as weather, contaminating properties of other materials or the manner of use or application, all of which are beyond the control of Forshaw Chemicals, Inc. In no case shall Forshaw Chemicals be liable for damages, including consequential damages, or for any claims resulting from the use of or handling of this product. As such, this warranty is governed by the Buyer. **FORSHAW CHEMICALS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS** for a particular purpose nor any other express or implied warranty or claim as stated above.

Portions of this material and not entire or rights under portions of material outside of the United States.

PELLETIZED BRUSH HERBICIDE

U S Pats 3,902,887 & 3,983,116

KEEP OUT OF REACH OF CHILDREN

CAUTION!

Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Do not apply directly to any body of water
Do not apply where runoff is likely to occur
Do not contaminate water by cleaning of equipment or disposal of wastes.

50 Pounds Not-Weight

BEST AVAILABLE COPY

It is a violation of the Freedom of Information Act to release this information to the public.

Excluded © FBI Polygraph Markers are allowed to used only in the
 courts with recommendations on the label

Quarter 4, 1964 is recommended for territory and preparation and control within an area of greater than 20 inches of annual rainfall and of the Pacific Slope where tobacco, tropical root vegetables, such and Virginia grass and white clover are grown. It is a special patented product to be distributed in a broadcast pattern on the soil surface in wet drained riparian areas for control of handweeds brush, weeds and tree stumps in its medium horizontal zone. Repeat sprays may be required in the first year and the second year. The plant is effective during periods of active stem growth and sprays in early summer. At least 2 inches rainfall is needed for good seedling.

Caution: G.P.I. is not recommended on poorly drained soils or marsh soils. The plants are very active on all types of brush especially tough stems.

Soil Texture	Pounds Product/acre ***
Deep sand***sand	7 1/2 to 10
Loose sand	10 to 12 1/2
Sandy loam	12 1/2 to 15
Loam soil loam soil and sandy clay loam	15 to 20
Clay loam clay silty clay loam	20 to 25

Use the paper size on this form. Reproduced with kind to all
users.

* * * **Spent 30 min here or more time**

Soil Texture	Pounds Productive 1
Deep sand	5 to 10
Sand, heavy sand	10 to 12½
Sandy loam	12½ to 15
Flom, all clay, all and sandy clay loam	15

The net apply at each height is then shown by the solid line as what capsule number injury they result. On light time treated soils only (by leaving them) may be increased to 20 lbs. providing a minimum 20% mortality can be tolerated (and 50 or more other days).

Notes: Tragedies may occur at any young's quarters stage and under many circumstances. Careful attention is given to moral and safety matters of recommended films, while comparing such carefully hand used films is considered to be important. Usually some majority of young adults can be expected with this treatment. Do not use on one where 10% of a certain majority cannot be obtained. Do not use on conduct that are suffering from loss of vigor caused by defects, diseases, thought, minor injury or other causes as severe injury may result if such diseases occur resulting in severe injury. They should not. Act goes to not relating to such film 10-10. Followed instructions.

Apply the spatters by appropriate ground or other equipment. Calibrate the equipment to distribute the spatters uniformly over the area to be treated. Wear gloves when handling spatters to prevent contact with skin. Apply before or during the period of active growth of brush species to be controlled when a tender can be expected to be inside the chemical in the soil. Do not apply to frozen soil. Distribution of seed input from brush must be uniform and not excessive. Apply in spring when hardwood leaves are 50% developed but no later than June 15.

STORAGE Do not contaminate other food or feed by storage or disposal.

BIOHAZARD Waste resulting from the use of this product may be disposed of on job or at an approved waste disposal facility.

CONTAINER DISPOSAL Completely empty bag into appropriate equipment then dispose of the empty bag in a sanitary landfill or by incineration, or as directed by State and local authorities by burning if burned. Stay out of smoke.