



When SPARTACIDE is used as directed, it will kill and prevent development of *Chlorella pyrenoidosa* (green), *Phormidium inundatum* (black algae), *Phormidium retzii* (Square D algae) *Lyngba versicolor* (blue-green) and other swimming pool algae. SPARTACIDE is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners, or premixed with other water-treating chemicals. SPARTACIDE works best when added by itself directly to pool water. SPARTACIDE at use dilution is non-corrosive to inanimate surfaces.

DIRECTIONS FOR USE:

INITIAL DOSE: 1 gallon SPARTACIDE to 50,000 gallons of water.

MAINTENANCE DOSE: 1 quart SPARTACIDE to 50,000 gallons of water.

Initial dose is used when filling pool at beginning of swimming season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or re-appear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance dose. Maintenance dose should be added at 3 - 5 day intervals. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently. If make-up water is added at the rate of more than 10% per week of pool capacity, add 1 quart of SPARTACIDE for each 10,000 gallons of fresh make-up water added per week.


ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides 8.4%
 n-Dialkyl (60% C14, 30% C16, 5% C12, 5% C18) methylbenzyl ammonium chlorides 1.6%

INERT INGREDIENTS:

90.0%
 100.0%

CONCENTRATED
 SWIMMING POOL
 ALGAECIDE



SPARTACIDE

26 WEST INMAN AVENUE,
 RAHWAY, N.J. 07065
 201-381-6577

DANGER. KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public waters. Apply this product only as specified on this label.

Rinse empty container thoroughly with water and discard.

NET CONTENTS:

E.P.A. Reg. No. 7245-5 E.P.A. Est. 7245-NJ-01