

CROWN

CHLORDANE 45 EMULSIFIABLE CONCENTRATE

FOLLOW DIRECTIONS CLOSELY. MEASURE ACCURATELY AND STIR EMULSION THOROUGHLY JUST BEFORE USING.

DILUTIONS

TO MAKE A 0.125% CHLORDANE EMULSION

FOR 0.25% CHLORDANE EMULSION

FOR 0.5% CHLORDANE EMULSION

FOR 0.75% CHLORDANE EMULSION

FOR 2% CHLORDANE EMULSION

HOW TO USE CROWN CHLORDANE 45

TICKS AND LICE ON HORSES

ANTS, ROACHES, FLIES, MOSQUITOES, GNATS, FLEAS, BROWN DOG TICKS, SILVERFISH, BOX ELDER BUGS, AND WATERBUGS INFESTING BARN, STABLES, SHEDS AND DOG KENNELS

FOR FURTHER INFORMATION, CONTACT YOUR LOCAL AGENT OR THE MANUFACTURER.

DIRECTIONS

FOLLOW DIRECTIONS CLOSELY. MEASURE ACCURATELY AND STIR EMULSION THOROUGHLY JUST BEFORE USING.

DILUTIONS

TO MAKE A 0.125% CHLORDANE EMULSION: Add 2 tablespoonsful Crown Chlordane 45 to 3 gallons of water; or 1 pint to 48 gallons; or 1 quart to 96 gallons.

FOR 0.25% CHLORDANE EMULSION: Add 4 tablespoonsful Crown Chlordane 45 to 3 gallons of water; or 1 pint to 24 gallons; or 1 quart to 48 gallons.

FOR 0.5% CHLORDANE EMULSION: Add ½ pint Chlordane 45 to 6 gallons of water; or 1 pint to 12 gallons; or 1 quart to 24 gallons.

FOR 0.75% CHLORDANE EMULSION: Add ½ pint Chlordane 45 to 4 gallons of water; or 1 pint to 8 gallons; or 1 quart to 16 gallons.

FOR 2% CHLORDANE EMULSION: Add ½ pint Chlordane 45 to 1½ gallons of water; or 1 pint to 3 gallons; or 1 quart to 6 gallons.

HOW TO USE CROWN CHLORDANE 45 after diluting with water for:

TICKS AND LICE ON HORSES: Spray thoroughly with 0.5% Chlordane Emulsion. For Spinose Ear Tick: Inject a small quantity of 0.5% Emulsion into the ears with syringe. Do not use on horses to be slaughtered for meat.

ANTS, ROACHES, FLIES, MOSQUITOES, GNATS, FLEAS, BROWN DOG TICKS, SILVERFISH, BOX ELDER BUGS, AND WATERBUGS INFESTING BARN, STABLES, SHEDS AND DOG KENNELS: Use 2% Chlordane Emulsion. Thoroughly wet all areas where these insects alight or crawl, particularly infested cracks and hiding places. For fleas and ticks, thoroughly and repeatedly spray infested areas of floors, and wall cracks as well as sleeping quarters of animals. Place fresh bedding in pet animal quarters after treatment.

ANTS, CHIGGERS, BROWN DOG TICKS AND SCORPIONS ON LAWNS: Spray lawn until wet with 0.25% or 0.5% Chlordane Emulsion. For a city lot 50 x 100 ft., use ½ to 1 pint Chlordane 45 in 12 or more

CROWN

CHLORDANE 45 EMULSIFIABLE CONCENTRATE

gallons of water. For field work 1 acre in 40 gallons of water or more diluted emulsion down entrance heavily around them for 5 to 10 feet off the treated area until they are washed into the soil and the birds are completely safe. Birds feeding over the area are not harmed.

CAUTION

Do not use in Poultry Houses, animal quarters after treatment of livestock, excepting horses. Do not feed forage crops treated with this product to animals or animals being finished for slaughter. Contact with skin can cause irritation. Wash hands thoroughly with soap and water after handling the concentrate or using the spray mist. Avoid breathing spray mist. Avoid contact with foodstuffs. Keep out of reach of children. Do not pour, store, or spill near heat or open flame. Do not use this product in the household. Do not use on fish and wildlife. Birds feeding over the treated area are not harmed. Keep out of any body of water when weather conditions favor runoff. Do not apply where runoff is likely to contaminate water by cleaning up or disposal of wastes. Use only as specified on label.

Do not reuse container. Destroy by crushing and burying in a safe place.


NOTICE OF WARRANTY

CROWN CHEMICALS, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE. SEE THE STATEMENTS ON THIS LABEL.

MANUFACTURED BY
CROWN CHEMICALS, INC.
ROCKFORD, ILL.



CHLORDANE 45 EMULSIFIABLE CONCENTRATE



gallons of water. For field work use 1 to 1½ quarts per acre in 40 gallons of water or more. Pour some of the diluted emulsion down entrance to ant hills, and spray heavily around them for 5 to 10 feet. Keep children and pets off the treated area until the insecticide has been washed into the soil and the treated area has dried completely. Birds feeding over treated lawn may be harmed.

CAUTION

Do not use in Poultry Houses. Place fresh bedding in animal quarters after treatment. **DO NOT USE ON DAIRY ANIMALS OR IN DAIRY BARN.** Do not apply to livestock, excepting horses. Do not apply to cats. Do not feed forage crops treated with this material to dairy animals or animals being finished for slaughter. Harmful if swallowed. Contact with skin can cause toxic symptoms. Wash hands thoroughly with soap and warm water after handling the concentrate or using the diluted product. Avoid breathing spray mist. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Do not use, pour, store, or spill near heat or open flame. Do not apply this product in the household. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Use only as specified on this label.

Do not reuse container. Destroy by perforating or crushing and burying in a safe place.

NOTICE OF WARRANTY

CROWN CHEMICALS, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statements on this label.

EPA REG NO 7273-29

MANUFACTURED BY
CROWN CHEMICALS, INC.
ROCKFORD, ILLINOIS

MEASURE
ROUGHLY

DN: Add 2
of water:

Add 4
of water:

1/2 pint
to 12

1/2 pint
to 8 gallons:

1/2 pint
to 3

er diluting

ughly with
k: Inject a
ears with
l for meat.

**GNATS,
SH, BOX
INFESTING
BEELES:** Use
areas where
treated cracks
roughly and
wall cracks
in bedding

KS AND
wet with
y lot 50 x
2 or more