



RAT AND MOUSE KILLER
with
DIPHACINONE

ACTIVE INGREDIENTS	BY WT.
Diphacinone (2-Diphenylacetyl-1,3-indandione)	0.005%
INERT INGREDIENTS	99.995%
TOTAL	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See additional cautions on back panel

DIRECTIONS

Place several self-feeding stations in locations frequented by rats and mice. Continuous feeding from 5 to 14 days should give control. Replace empty stations within 24 hours. Bait should be available to rats for at least 15 days or until complete lack of feeding is noticed. Wet or otherwise decomposed bait should be replaced immediately.

Mice use at least four ounces of bait per station. Space stations more than 15 feet apart. (preferable 8-12 feet apart)

Rats use at least a half pound of bait per station, increasing the amount if heavy infestation exists.

Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.

CAUTION

Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of 2-hydroxycoumarin. Bait should be placed in locations not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

Do not re-use container. Destroy when empty.

Manufactured by
CROWN CHEMICALS, INC.
Rockford, Illinois

U.S.D.A. Reg. No. 7273-44

Net Contents 5 lbs