


DIRECTIONS

Vegetables in Home Gardens

For Mexican Bean Beetles, Leafhoppers, Lygus and Stink Bugs on beans, apply dust to uniformly cover upper and lower leaf surfaces. Dust every 7 to 10 days or as needed. May be applied up to harvest at rate directed without exceeding residue tolerances.

For Striped and Spotted Cucumber Beetles, Squash Bugs, Flea Beetles, Pickleworm and Melonworm on cucumbers, summer squash dust to uniformly cover upper and lower leaf surfaces. Dust every 7 to 10 days or as needed. May be applied up to harvest at rate directed without exceeding residue tolerances.

For Tomato Fruitworm, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer and Stinkbugs on tomatoes, apply dust to uniformly cover upper and lower leaf surfaces. Dust every 7 to 10 days or as needed. May be applied up to day of harvest at rate




POLTRY DUST

ACCEPTED
1975

CAUTION

**KEEP OUT OF REACH OF
CHILDREN AND ANIMALS**

Harmful if swallowed, inhaled, or absorbed through skin. Do not take internally. Avoid breathing dust. Avoid contact with skin. Avoid unnecessary contact. Wash thoroughly after handling and before eating. Take shower after work. Change contaminated clothing daily. Wear regular long sleeved clothing. Avoid contamination of feed and foodstuffs. Do not feed treated vines or trash to meat or dairy animals. Keep



THE NEW YORK PUBLIC LIBRARY
ASTOR LENOX TILDEN FOUNDATION
500 5TH AVENUE
NEW YORK 17, N.Y.