

Dec 4, 1969

7273-55

original, it for the use there is reserved when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential or punitive damages such as loss of profits or value, etc.

CROWN CHEMICALS, INC. makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY.

BUYER assumes the risk of any, and all, damage to the contents of any order at normal and through order conditions for assembly, for installation, for use and for maintenance.

HOUSEFLIES

Robert W. Long, Jr. For the control of hepatitis A virus, the first thing you need to know is the symptoms of the virus. The symptoms of hepatitis A are usually mild and last for about two to four weeks. The symptoms include fatigue, loss of appetite, nausea, vomiting, and jaundice. The virus is spread through contaminated food and water. The best way to prevent hepatitis A is to avoid contaminated food and water. If you are traveling to a country where hepatitis A is common, you should get vaccinated. The vaccine is safe and effective. It is also important to practice good hygiene, such as washing your hands thoroughly with soap and water. If you have been exposed to the virus, you should see your doctor for a blood test. The test will tell you if you have the virus. If you do, you should rest and drink plenty of fluids. You should also avoid alcohol and fatty foods. The virus will usually go away on its own. In most cases, the symptoms will disappear within a few weeks. However, in some cases, the virus can cause long-term damage to the liver. This is rare, but it can happen. If you have any concerns about hepatitis A, you should talk to your doctor. They can help you understand the virus and how to prevent it. They can also help you if you have been exposed to the virus. Remember, hepatitis A is a preventable disease. By taking the right steps, you can protect yourself and your family. Stay healthy and safe!

Figure 1. The effect of the concentration of the Fe^{2+} solution on the adsorption of Fe^{2+} by the Fe^{2+} -loaded adsorbent. The concentration of the Fe^{2+} solution was 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 50, 100, 200, 500, 1000, 2000, 5000, 10000, 20000, 50000, 100000, 200000, 500000, 1000000, 2000000, 5000000, 10000000, 20000000, 50000000, 100000000, 200000000, 500000000, 1000000000, 2000000000, 5000000000, 10000000000, 20000000000, 50000000000, 100000000000, 200000000000, 500000000000, 1000000000000, 2000000000000, 5000000000000, 10000000000000, 20000000000000, 50000000000000, 100000000000000, 200000000000000, 500000000000000, 1000000000000000, 2000000000000000, 5000000000000000, 10000000000000000, 20000000000000000, 50000000000000000, 100000000000000000, 200000000000000000, 500000000000000000, 1000000000000000000, 2000000000000000000, 5000000000000000000, 10000000000000000000, 20000000000000000000, 50000000000000000000, 100000000000000000000, 200000000000000000000, 500000000000000000000, 1000000000000000000000, 2000000000000000000000, 5000000000000000000000, 10000000000000000000000, 20000000000000000000000, 50000000000000000000000, 100000000000000000000000, 200000000000000000000000, 500000000000000000000000, 1000000000000000000000000, 2000000000000000000000000, 5000000000000000000000000, 10000000000000000000000000, 20000000000000000000000000, 50000000000000000000000000, 100000000000000000000000000, 200000000000000000000000000, 500000000000000000000000000, 1000000000000000000000000000, 2000000000000000000000000000, 5000000000000000000000000000, 10000000000000000000000000000, 20000000000000000000000000000, 50000000000000000000000000000, 100000000000000000000000000000, 200000000000000000000000000000, 500000000000000000000000000000, 1000000000000000000000000000000, 2000000000000000000000000000000, 5000000000000000000000000000000, 10000000000000000000000000000000, 20000000000000000000000000000000, 50000000000000000000000000000000, 100000000000000000000000000000000, 200000000000000000000000000000000, 500000000000000000000000000000000, 1000000000000000000000000000000000, 2000000000000000000000000000000000, 5000000000000000000000000000000000, 10000000000000000000000000000000000, 20000000000000000000000000000000000, 50000000000000000000000000000000000, 100000000000000000000000000000000000, 200000000000000000000000000000000000, 500000000000000000000000000000000000, 1000000000000000000000000000000000000, 2000000000000000000000000000000000000, 5000000000000000000000000000000000000, 10000000000000000000000000000000000000, 20000000000000000000000000000000000000, 50000000000000000000000000000000000000, 100000000000000000000000000000000000000, 200000000000000000000000000000000000000, 500000000000000000000000000000000000000, 1000000000000000000000000000000000000000, 2000000000000000000000000000000000000000, 5000000000000000000000000000000000000000, 100, 200, 500, 1000, 2000, 5000, 100, 200, 500, 1000, 2000, 5000, 100, 200, 500, 1000, 2000, 5000, 100, 200, 500, 1000, 2000, 5000, 100, 200, 500, 100000000

CONSIDER A CERTAIN FLOCK AS A RESIDUAL, THE NEXT YEAR, AND AGAIN IN THE FOLLOWING YEAR MEAT PLANTS IN THE AREA MEAT PLANTS ARE PRODUCED AND HANDLED.

1. *Journal of the American Medical Association*, 1997; 278: 1039-1044.

1. NAME
 2. ADDRESS
 3. PHONE

CYGN* 2-E

Systemic Inequality
The concept of income and other
in organized groups,
refer to attached exhibit.

ACTIVE PERSONS

the 1990s, the number of people in the world who are under 15 years of age is expected to increase from 1.1 billion to 1.5 billion. The number of people aged 65 and over is expected to increase from 200 million to 400 million. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion.

1. *Journal of the American Medical Association*, 2000; 283: 2689-2696.

WARNING. KEEP OUT OF REACH OF CHILDREN

See side panel for antidote and other cautions.

NET CONTENTS: 1 Quart

CAUTION!

VAPOR HARMFUL HARMFUL OR FATAL IF SWALLOWED CONCENTRATED MATERIAL CAUSES EYE IRRITATION COMBUSTIBLE

August 1953, 1000 ft. or 1000 ft. or 1000 ft.

August 1964

On 21, 22 and 23 June 1992, the water level

Vibrio anguillarum, *Vibrio parvulus*

1. What is the purpose of the study?

As noted above, the results of the posttest are quite striking. The mean score on the posttest was 10.8, with a standard deviation of 1.8. This is a significant improvement over the mean score of 7.5 on the pretest, with a standard deviation of 2.0. The results of the posttest are also consistent with the results of the pretest, in that the mean score on the posttest was significantly higher than the mean score on the pretest.

NOTE: PHYLIP 3.42 is used to determine phylogeny of the 16S rDNA sequences. The bootstrap values are indicated.

The following table shows the results of the regression analysis. The dependent variable is the number of days of absence from work due to illness. The independent variables are the age, sex, and education of the respondent, and the number of children in the household. The results show that the number of days of absence from work due to illness increases with age, and is higher for females than for males. Education has a negative effect on the number of days of absence from work due to illness, and the number of children in the household has a positive effect.

[illegible][illegible]

DIRECTIONS FOR USE
DISCLAIMER

CYGON^{*} 2-E

Systemic Insecticide
CONTROLS HOUSEFLIES
For control of Insects and Mites
on ornamental plants,
refer to attached leaflet

WARNING: KEEP OUT OF REACH OF CHILDREN

See label and directions for use on other insects.

© 1970 FMC CORPORATION