

DIRECTIONS FOR USE

HOUSEHOLD INSECTS

COCKROACHES (WATERBUGS), ANTS, AND SILVER FISH IN BUILDINGS: (By Pest Control Operators Only) For control of ants, silverfish, and resistant cockroaches (waterbugs), prepare a 2% spray by thoroughly mixing 2 fl. ozs. (5 tablespoons) of this product with sufficient water to make one gal. of finished spray and make spot applications of this mixture to areas infested with these insects. Application should be made as a coarse spray & with a paint brush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves and similar areas where these insects hide. Treatments should be repeated as necessary. Under conditions of heavy cockroach infestation, it is suggested that the first application be made at a strength of 1% this product (5 fl. ozs. this product per gal.), followed by monthly applications of 1/2% this product (2 1/2 fl. ozs. this product per gal.).

BROWN DOG TICKS IN BUILDINGS: Prepare a 2% this product spray by thoroughly mixing 2 fl. ozs. (5 tablespoons) of this product with sufficient water to make 1 gal. of finished spray. Thoroughly spray infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets and localized areas of floors and floor coverings. Applications should be repeated as infestations warrant and as reinfestations occur. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

When used in dwellings care should be taken to avoid deposits which could be frequently contacted by children. Care should also be taken to avoid spilling of wallpaper and fabrics. Do not contaminate food, food containers or cooking utensils.

ORNAMENTAL TURF AND ORNAMENTAL INSECTS
ORNAMENTALS: For control of those insects listed below attacking arborescences, azaleas, carnations, chrysanthemums, holly, juniper, rhododendron, roses, pine, spruce, fir, and other flowers, shrubs and shade trees mix 1 fl. oz. (2 tablespoons) in 3 gals. of water and apply as a thorough foliage spray using a knapsack or hand type spray. Treatment should be made when insects first appear and repeated as infestations warrant.

Aphids	Cyclops mites
Scale insects	Caterpillars
Spider mites	Fire beetles
Carpet beetles	European spruce sawfly
Thrips	Two Spotted mite
Holly Bud Moths	Scale crawlers (Cottin)
Dipterous Leaf Miners	European spruce sawfly
Leafhoppers	North American spruce sawfly
Whiteflies	Starlings

NOTE: Do not use on plants which are sensitive to diazinon. Do not use on plants which are sensitive to diazinon.

EPA Reg. No. 7273-65 AA
(P. D. No. 523 IL 1)

CROWN

DZ-25

(25% EMULSIFIABLE DIAZINON)

INSECTICIDE



KEEP OUT

Do not use on plants which are sensitive to diazinon. Do not use on plants which are sensitive to diazinon. Do not use on plants which are sensitive to diazinon.

DO NOT USE OR STORE IN PLACES WHERE CHILDREN CAN GET AT IT

MANUFACTURED BY
CROWN CHEMICALS

ROCKFORD, ILLINOIS 61103 USA

NET CONTENTS

8.6 GALLON

CROWN

DZ-25

(25% EMULSIFIABLE DIAZINON)

INSECTICIDE

POISONED CHILD ONLY

POISONED CHILD ONLY

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

**MANUFACTURED FOR
CROWN CHEMICALS
ROCKFORD, ILLINOIS 61103 USA**

NET CONTENTS 85 GALLON

DIRECTIONS FOR USE

FLEAS, TICKS AND CHIGGERS INFESTING LAWNS AND OTHER OUTDOOR AREAS: Use 12 fl. ozs. per 5 gals. of water and apply at the rate of 1 gal. of spray per 1,000 sq. ft. of infested area.

LAWN CHINCH BUGS: For control of chinch bugs in lawns, thoroughly wet down the grass and apply this product at the rate of 4 to 6 fl. ozs. in 25 gals. of water per 1,000 sq. ft. Applications should be repeated in 7-10 days if necessary.

FLY CONTROL ON FARM AREAS AND BUILDINGS: Do not use in milk rooms, dairy barns or poultry houses.

RESIDUAL SPRAYS: Apply a spray prepared by mixing 2 qts. of this product in 25 gals. of water as an overall application to the ceilings and walls of livestock sheds, calf barns, hog barns and other farm buildings. Other areas where flies congregate, such as around garbage dumps and corrals, should be sprayed thoroughly. Repeat applications as fly populations warrant.

Longer residual control may be obtained by increasing the dosage to 4 qts. of this product in 25 gals. of water. Applications should be made to the point of runoff. One gal. of the above spray should cover from 350 to 750 sq. ft. of area depending on surface to be treated.

NOTE: Remove animals from buildings during spraying operations and keep them out for at least 4 hours. Do not spray animals or contaminate their feed or drinking water. Do not use in poultry houses. Do not use in dwellings, milk houses, or dairy barns.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift or spray from areas treated.

Do not contaminate water by cleaning of equipment or disposal of waste.

NOTICE OF WARRANTY

CROWN CHEMICALS, INC. warrants that the product is free from defects in material and workmanship under normal use and conditions for a period of 90 days from the date of purchase.

CROWN CHEMICALS, INC. warrants that the product is free from defects in material and workmanship under normal use and conditions for a period of 90 days from the date of purchase.

**CHECK
PEST**

ACCEPTED
7277-17



INSE

**CHECK
PEST**

May be fatal if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not get in eyes, on skin or on clothing. Do not breathe dust. Wear protective clothing. Wash all contaminated clothing with soap and hot water before reuse. Do not contaminate feed, feeders or feedstuffs. If spilled, discontinue handling immediately and discontinue use. Atropine is a sedative. Do not use it in its weak form feed or feed supplements in a young.

The present study was designed to assess the effect of a 12-week intervention on health-related quality of life in patients with chronic low back pain, and to determine the impact of compliance. A pilot study was conducted first.

11 12 13 14



INSECTICIDE 106

(FOR FOGGING USE ONLY BY
TRAINED PERSONNEL)

