



**DIRECTIONS FOR USE
IN ANY TYPE SPRAYER**

For use in non-crop areas such as fence rows, industrial sites, storage yards and rights of way.

FOR NOXIOUS WEEDS such as annual saw thistle, beggar ticks, bitterweed, bull thistle, burdock, cocklebur, dandelion, fanweed, heal-all, hedge nettle, henbit, Jimson weed, lambs quarters, pennywort, milk purslane, mustards, pigweeds, plantains, poison hemlock, puncture vine, purslane, ragweed, wild licorice, wild radish and star thistle. Apply when weeds are growing vigorously.

Dilute at the rate of one part Weed & Brushoff to 100 parts of water and apply as a coarse spray.

For small areas, mix one teaspoon of Weed & Brushoff to one quart of water and apply as a coarse spray to cover about 50 square feet. For large areas, mix one gallon Weed & Brushoff in 100 gallons of water. Coverage, about one acre or 40,000 sq ft. Injury may occur when this product is used at these rates on Carpet, St Augustine, Bermuda or other creeping grasses. Use only on established grasses.

FOR WOODY WEEDS AND BRUSH poison ivy, sumac, honeysuckle, poison oak, brambles, birch, buckbrush, salt cedar, chokecherry, wild cherry, cottonwood, box elder, mulberry, osage orange, poplar, raspberry, cherokee rose, sagebrush, trumpet vine, willow and witch hazel. Mix at the rate of one part Weed & Brushoff to 50 parts of water and apply as a coarse spray and wet the plants thoroughly when in full leaf.

For small areas, mix one part Weed & Brushoff in 10 gallons of water and for large areas, mix one gallon Weed & Brushoff in 50 gallons of water. Spot spray after six weeks if necessary.

FOR STUMP TREATMENT: Mix one gallon Weed & Brushoff in 20 gallons of diesel or fuel oil and spray thoroughly over tops and sides of stump at any time.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.



WEED AND BRUSH **OFF**

ACCEPTED

JAN 11 1972

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
REGISTRATION NUMBER REGISTERED
122385



BRUSHOFF KILLS

Sow Thistle (Annual)	Pennywort	Honeysuckle
Beggar Ticks	Milk Purslane	Osage Orange
Bitterweed	Mustards	Poison Oak
Bull Thistle	Pigweeds	Poplar
Burdock	Plantain	Brambles
Cocklebur	Poison Hemlock	Raspberry
Dandelion	Puncture Vine	Birch
Fanweed	Purslane	Cherokee Rose
Heal-all	Ragweed	Buckbrush
Hedge Nettle	Wild Licorice	Sagebrush
Henbit	Wild Radish	Salt Cedar
Jimson Weed	Thistle	Trumpet Vine
Lambs Quarter	Chickweed	Choke Cherry
Poison Ivy	Box Elder	Willow
Sumac	Mulberry	Witch Hazel
Cottonwood		

CAUTION: Harmful if swallowed.

May cause skin irritation. Avoid contact with skin, eyes or clothing.

Avoid a dry drift to susceptible plants as this product may injure other plants near spray drift or over ornamentals etc. Coarse spray is most likely to drift. Do not use same sprayer for other purposes.

Do not store near fertilizers, seeds, insecticides, fungicides or do not use in greenhouse. Do not use around the home, recreation areas or similar areas.

Do not contaminate water used for irrigation or domestic purposes.

Do not reuse containers. Destroy by puncturing, crushing or burying in a well-drained area from water or domestic vegetation. Do not burn.

Do not spray on open soil or near open flame.

Crown Chemicals Division A Division of International Minerals and Chemical Company, otherwise express or implied, constitutes the sole and exclusive warranty extended beyond the period stated above.