

## DIRECTIONS

Direction should be followed closely. Proper application is essential to the successful use of this product. There are four general methods for applying this product:

### FOLIAGE APPLICATION

Mix 3 quarts of this product in 100 gallons of water and spray thoroughly all foliage and stems to ground line when plants are in full leaf and up until 2 weeks before expected frost date in your area.

When resistant brush such as oak, maple, persimmon, hickory and certain oaks constitute only a small proportion of mixed brush, use 4 quarts of this product in 100 gallons of water. Where these plants constitute a major part of the brush stand follow the procedure described under stump or basal treatment.

### Tree STUMP TREATMENT

(For use on cut stumps over 4" in diameter. On woody plants under 4" in diameter, use Foliage Application and retreat resistant species using basal spray.)

Mix 1 gallon of this product in 50 gallons of Diesel oil or Kerosene and apply with a low volume knapsack sprayer using solid cone nozzle of medium orifice. Completely wet top and sides of stump to ground level and spray all exposed lateral roots. Treatment can be made immediately following cutting or any time during the year.

For the more resistant woody plants as named above use 1 gallon of this product in 25 gallons of oil.

### BASAL SPRAY

(Summer or Dormant)

This treatment is used for the harder-to-kill woody plants and consists of applying this product in Diesel oil or Kerosene to the basal part of standing brush at any time of the year. Use 1 to 2 gallons this product in 50 gallons oil and apply with low volume knapsack or power equipment. Apply material to all sides of stems from ground line up 12 to 18 inches. Use enough material to cause run-off. Use most concentrated solution on oak, basswood, persimmon, beech, oak and red maple. For dormant spray the closer the material is applied to bud break the more effective in suppressing regrowth.

### FRILL METHOD

(For use on large trees)

Make cut with axe around trunk of the tree at waist height. Pour a solution of 1 gallon this product in 50 gallons water in the open cut. Pour in as much as frill will hold without wasting. Use 1 gallon this product in 25 gallons water on more resistant varieties such as persimmon, beech, hickory and sweet gum.

NET CONTENTS  
GALLONS

GEN

This product is a combination of 2,4-D emulsifiers added.

This product mixes with water. It also is suitable for plane applications.

This product is an

- Aspen
- Buck
- Blackberry
- Box Elder
- Buckbrush
- Butternut
- Catalpa
- Chestnut
- Craapple
- Ground Cherry
- Dogwood
- Dawberries

on utilization of problem. Tolerant of control.

This product can be uncropped land, but avoid in crops.

When diluting, the material should be stirred up.

Harmful if swallowed.

Avoid contact with

Wash contaminated


Avoid spray drift to cotton, beans, peas, drift). Do not use or store near fertilizers, product may injure or contaminate water or greenhouse. Do not use in similar areas.

Do not graze dairy application.

Do not graze meat or

This product will kill contaminate water. Destroy containers. Do not burn.

Crown Chemicals for purpose, or otherwise use, which extend to



# LOW VOLATILE BRUSH & WEED KILLER

Contains

2 2-3 Lbs. 2,4-D and 1 1-3 Lbs. 2,4,5-T

ACID EQUIVALENT PER GALLON

### ACTIVE INGREDIENTS:

- \*Iso-Octyl Ester of 2,4-Dichlorophenoxyacetic Acid.....42.4 %
- \*\*Iso-Octyl Ester of 2,4,5-Trichlorophenoxyacetic Acid.....20.94 %

### INERT INGREDIENTS

- \*Equivalent to 28.12% by weight of 2,4-Dichlorophenoxyacetic Acid.
- \*\*Equivalent to 14.55% by weight of 2,4,5-Trichlorophenoxyacetic Acid.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

See side panel for other warnings.

Manufactured by


**CROWN CHEMICALS**

ROCKFORD, ILL. 61103

77777

NET CONTENTS  
GALLONS

GENERAL INSTRUCTIONS



ACCEPTED  
Aug 9, 1971  
FEDERAL AGRICULTURAL MECHANICAL INSECTICIDES  
REGISTRATION DIVISION / CT  
EPA REG. NO. 7273-88

*Comment*

# LOW VOLATILE BRUSH & WEED KILLER

Contains

2 2-3 Lbs. 2,4-D and 1 1-3 Lbs. 2,4,5-T  
ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS:

- \*Iso-Octyl Ester of 2,4-Dichlorophenoxyacetic Acid ..... 42.4 %
- \*\*Iso-Octyl Ester of 2,4,5-Trichlorophenoxyacetic Acid ..... 20.94 %

INERT INGREDIENTS

- ..... 36.66 %
- \*Equivalent to 28.12% by weight of 2,4-Dichlorophenoxyacetic Acid.
- \*\*Equivalent to 14.55% by weight of 2,4,5-Trichlorophenoxyacetic Acid.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

See side panel for other warnings.

Manufactured by

## CROWN CHEMICALS

ROCKFORD, ILL. 61103

This product is a low volatile, powerful, economical combination of 2,4-D and 2,4,5-T esters — penetrant, and emulsifiers added.

This product mixes with oil or kerosene and emulsifies in water. It also combines with oil water mixtures for air plane applications.

This product is an effective formulation for controlling

Aspen	Elderberry	Spicebrush
Birch	Elm	Sunoc
Blackberry	Honeysuckle	Willow
Box Elder	Horse Chestnut	Sweet Gum
Buckbrush	Locust	Sycamore
Butternut	Oak	Walnut
Catalpa	Csage	Wild Cherry
Chestnut	Poison Ivy	Wild Grape
Crabapple	Poison Oak	Wild Rose
Ground Cherry	Poplar	Virginia Creeper
Dogwood	Raspberry	and many other species.
Dewberries	Sassafras	

on utility ~~road~~ ~~along~~ ~~highways~~ or wherever brush is a problem. Tolerant species may require repeat applications for effective control.

This prod. can be used to control many noxious perennial weeds on uncrowded land, but is not recommended for selective treatment of weeds in crops.

When diluting, the material must be mixed thoroughly before using. If the material is allowed to stand for a length of time after dilution, it should be stirred again before using.

### CAUTION

Harmful if swallowed.

Avoid contact with eyes, skin or clothing. May cause skin irritation.

Wash contaminated clothing before re-use.

### WARNING

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift). Do not use same spray equipment for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation purposes. Do not use in a house. Do not use in or around the home, recreational sites or other areas.

Do not graze dairy animals on treated areas within 6 weeks after application.

Do not graze meat animals on treated areas within 2 weeks of slaughter.

This product will kill fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Destroy containers by puncturing, crushing and burying in a safe area. Do not burn.

### NOTICE OF WARRANTY

Crown Chemicals makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied concerning this product or its use, which extend beyond the statements on this label.

USDA Reg No 7273-88

