

NET GALLONS

DIRECTIONS FOR USE

CUTWORMS AND ARMYWORMS

Treat cutworms and Armyworms in grain crops such as wheat, rye or barley, and corn. Use 1 to 2 quarts of this product in 10 to 20 gallons of water per acre. Do not feed treated fodder or straw to dairy animals or animals being finished for slaughter or allow these animals to graze treated areas.

GRASSHOPPERS

TOXAPHENE insecticides can be used to advantage on infestations of young grasshoppers found on roadsides, field margins, or idle land next to cultivated fields. Spraying these infestations when the main hatch is completed or when the young hoppers begin to move off the hatching grounds greatly reduces the acreage to be treated. Use 1 to 1 1/3 quarts of this product in 10 to 20 gallons of water per acre.

TOXAPHENE also prevents grasshoppers from migrating such crops as corn, sorghum, cotton and soybeans. If field margins are treated before the hoppers move in. The application of from 2/3 to 1 quart of this product in 10 to 20 gallons of water per acre, when used as a spray, depending on the height and density of vegetation, is recommended for general use. A one-third increase in the dosage of spray may be necessary late in the season, when most of the grasshoppers are adults and vegetation is maturing.

SPECIAL NOTES


Do not treat sorghum field margins within 28 days of harvest or soybean field margins within 21 days of harvest. Do not treat sorghum more than once after heads start to form. Do not feed treated soybean forage to poultry. Do not feed treated soybean, fish feed, to poultry, dairy animals, or animals being finished for slaughter.

Do not feed treated forage, straw, or cotton trash to dairy animals or animals being finished for slaughter. Do not store products from storage temperature below 15°F.

EXTREMELY TOXIC to fish and wildlife. DO NOT allow this material, or any body of water, to be applied to any area as specified in the label. Do not apply when weather conditions favor drift from areas treated. Birds feeding on treated areas may be killed. Do not apply when drift is likely to occur.

Do not contaminate water by cleaning of equipment or disposal of waste.

Do not use containers. Dispose of container by puncturing, then salting or by crushing and burying in a suitable area away from water supply.



T-6 TOXAPHENE

(Contains 6 lbs. TOXAPHENE per gal. One qt. contains 1.5 lbs. Toxaphene)

AN AGRICULTURAL INSECTICIDE

WARNING: KEEP OUT OF REACH OF CHILDREN

Extremely hazardous if swallowed, inhaled or absorbed through skin or eyes! DO NOT breathe spray mist. Wear a suitable respirator. DO NOT get in eyes, on skin or on clothing. Wear goggles. DO NOT leave in eyes, on skin or clothing. Wash thoroughly with soap and water. AVOID contact with food and food stuffs. KEEP AWAY from domestic animals. Not for use on or around the house.

ACTIVE INGREDIENTS

Toxaphene (Technical Chlorinated Camphene) 60.17
Containing 67.69 Chlorine 36.91
Light Petroleum Distillate 2.92

INERT INGREDIENTS

2.99
100.00

MANUFACTURED BY

CROWN CHEMICALS

ROCKFORD, ILL. 61101

MIXING INSTRUCTIONS

Carefully pour required amount of concentrate into nearly filled spray tank with agitator. Another method is to pour concentrate into other suitable container. Add an equal volume of water and stir thoroughly with paddle until creamy white. Carefully pour the cream into nearly filled spray tank. Then add water to tank. Run agitator for several minutes before spraying.

CAUTION: If concentrate does not form a suspension with water but tends to separate into a layer, do not use it as injury to plants may result.

RECOMMENDATIONS FOR SPRAYING AGRICULTURAL CROPS

DO NOT USE WITHOUT DILUTION, mixing as directed in the proper amount per acre completely average with the type of nozzle used. For general spraying use 10 to 20 gallons of water and 1 to 2 quarts of product per acre. For spot spraying use 3 to 10 gallons of water and 1/2 to 1 quart of product per acre. ALFALFA LEECH, LEECH BUGS, SOYBEAN SEED, Apply 1 quart of this product per acre.

BEAN LEAF MINER on beans. Apply twice set and an acre inches average for beans. Make two applications at 15 day intervals, apply after pods begin to form. Apply 1 quart of product per acre.

COTTON INSECT CONTROL

Leafworm, Budworm, Rapid Plant Borer, Leafworm, Tarnished Plant Bug, Cotton Bollworm start control as soon as egg hatched worms are found.

For Cotton Leafworm apply when worms appear on leaves and spray with 2 quarts of this product per acre.

SERPENTINE LEAF MINER

on Soybean. For Soybean apply 2 quarts per acre of this product. Do not apply after start of pod formation.

CRISPER BUG, CRISPER MINER on alfalfa. Apply 1 quart of this product per acre. Use at first growth in seed row.


For further information contact your local agricultural extension office.

NOTICE OF WARRANTY

CROWN CHEMICALS makes no warranty of merchantability, fitness for any purpose, or express or implied concerning the product or extends beyond the state in which sold.

EPA REG.

NET GALLONS



T-6 TOXAPHENE

(Contains 6 lbs. TOXAPHENE per gal. One qt. contains 1.5 lbs. Toxaphene)

AN AGRICULTURAL INSECTICIDE

WARNING: KEEP OUT OF REACH OF CHILDREN

Extremely hazardous if swallowed, inhaled or absorbed through skin or eyes. **DO NOT** breathe spray mist. Wear a suitable respirator. **DO NOT** get in eyes, on skin or on clothing. Wear goggles. **DO NOT** get in eyes, on skin or clothing. Wash thoroughly with soap and water. **AVOID** contamination of food and food stuffs. **KEEP AWAY** from domestic animals. Not for use on or around the home.

ACTIVE INGREDIENTS

Toxaphene (Technical Chlorinated Camphene)	50.17
Containing 67-69% Chlorine	36.83
Light Petroleum Distillate	2.02

INERT INGREDIENTS

100.00

MIXING INSTRUCTIONS

Carefully pour required amount of T-6 insect into partially filled spray tank with agitator running. Another method is to pour T-6 insect into bucket or other suitable container. Add an equal amount of water and stir thoroughly with paddle until mixture is creamy white. Carefully pour the creamy mixture into nearly filled spray tank. Then add water to fill tank. Run agitator for several minutes before spraying.

CAUTION: If concentrate does not form milky suspension with water but tends to separate as an oily layer, do not use it as injury to plants may result.

RECOMMENDATIONS FOR SPRAYING AGRICULTURAL CROPS

DO NOT USE WITHOUT DILUTION. Dilute by mixing as directed in the proper amount of water to give complete coverage with the type of equipment used. In ground sprayers use 10 to 20 gallons of water, and by airplane use 3 to 10 gallons of water per acre, depending upon growth of crop.

ALFALFA WEEVIL, LYBUS BUGS - On alfalfa seed. Apply 1 quart of this product per acre as needed.

BEAN LEAF MINER - Beans - Apply after beans have set and whenever miners average four to a leaf. Make two applications at 15 day intervals. Do not apply after pods begin to form. Apply 1 quart of the product per acre.

COTTON INSECT CONTROL

Boll Weevil, Bollworm, Rapid Plant Bug, Cotton Leafworm, Tarnished Plant Bug, Cotton Fleahopper. For Bollworm start control as soon as eggs and newly hatched worms are found.

For Cotton Fleahopper apply when cotton first appears, does not need to grow. Apply 2 1/2 pints of this product per acre.

SERPENTINE LEAF MINER

Corn, Beans, Tomatoes, Etc. Rate per acre: 1 quart of the product. Do not apply after fruit or seed parts of the crop start to form.

SPITTLEBUG, WIMPHS - Seed, Alfalfa, Etc. Apply 1 quart of this product per acre. The only one fast acting seed crops.

For further specific information consult the representative extension service.

NOTICE OF WARRANTY

CROWN CHEMICALS makes no warranty of merchantability, fitness for any purpose or otherwise, express or implied, concerning this product or its use which extend beyond the statements on this label.

CROWN CHEMICALS

ROCKFORD, ILL. 61103

EPA REG. NO. 7275-6