

DIRECTIONS

DAIRY BARNS, MILK SHEDS, STABLES, LIVE STOCK BARNS, LOAFING SHEDS, PIG PENS, AND OUTDOOR AREAS: Scatter dry bait at the rate of $\frac{1}{4}$ lb. per 1,000 sq. ft. on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways, and window sills. On freshly lined floors, place treatment on feed sacks or on paper. Repeat procedure as necessary for practical fly control.

CAGED POULTRY AND TURKEYS ON WIRE: Scatter dry bait at the rate of $\frac{1}{4}$ lb. per 100 sq. ft. on droppings beneath cages or porches. Repeat at weekly intervals.

POULTRY HOUSES: Scatter "Malvex" dry bait at the rate of $\frac{1}{4}$ lb. bait per 1,000 sq. ft. of area where flies congregate, but keep out of reach of birds. Apply to manure, on window sills, outside of penned enclosures and on floor of feed storage room. Repeat as necessary.

DOG KENNELS: Scatter "Malvex" dry bait at the rate of $\frac{1}{4}$ lb. per 1,000 sq. ft. outside dog runways, on window sills, ledges and other surfaces out of reach of animals. For caged units, apply to dropping area. Repeat as necessary.

FOOD PROCESSING PLANTS: For control of resistant or non-resistant houseflies around the outside premises of food processing plants, use $\frac{1}{4}$ lb. per 500 square feet of fly breeding area. Scatter on areas where flies congregate.

GARBAGE CANS: Apply bait around and inside garbage can. Repeat as necessary.

FOR REDUCING POPULATIONS OF RESISTANT AND NON-RESISTANT HOUSEFLIES IN AND AROUND OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREAS, PATIOS, LOADING DOCKS, AND OUTDOOR LATRINES, PARKING AREAS, REFUSE AREAS AROUND SERVICE STATIONS, OPEN AIR DRIVE-INS, ICE CREAM STANDS, GARBAGE COLLECTION AND DISPOSAL AREAS: Use "Malvex" dry bait at $\frac{1}{4}$ lb. per 1,000 sq. ft. of area. Apply on ground or areas where flies congregate. For open air drive-ins, ice cream stands and restaurants, apply on apron between serving counter and parking area, paying particular attention to trash barrels and garbage cans where flies congregate.

CROWN

'MALVEX'

DRY FLY BAIT

FOR FARM AND OUTSIDE HOME USE

For Active Ingredients, see back panel.
For Complete List of Ingredients, see back panel.
For Full Directions, see back panel.
For Full First Aid, see back panel.

ACTIVE INGREDIENTS		BY WT.
2,2-Dichloro-4,4'-dimethyl		46%
phosphate		
Residual Carcinogen		.04%
INERT INGREDIENTS		99.50%
TOTAL		100.00%

*Equivalent to 0.5% of Vapona® Insecticide
® Trademark of Shell Chemical Co.

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL PRECAUTIONS ON BACK PANEL

DO NOT REUSE EMPTY CONTAINER. DESTROY
WHEN EMPTY.

NET WEIGHT 1 LB.

Do not contaminate milk, milk handling equipment, feed or drinking water.

7273-105

CAUTION

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs. Keep out of reach of children. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidote.

This product is toxic to fish, wildlife and birds. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE OF WARRANTY

CROWN CHEMICALS, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statements on this label.

Manufactured for
CROWN CHEMICALS
ROCKFORD, ILLINOIS 61101

EPA Reg. No. 7273-105
EPA EST. No. 5-3-11-1