

No. HVX-1

NET WT. 50 LBS.



March 29, 1972

7272-107

TERRA-VAR

GRANULAR WEED AND BRUSH KILLER

ACTIVE INGREDIENTS:

Bromacil
(5-Bromo-3-sec-butyl-6-methyluracil) 2.00%

INERT INGREDIENTS 98.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN. May cause irritation of eyes, nose, throat and skin.
Avoid breathing dust. Avoid contact with skin, eyes, and clothing.

Do not reuse container. Destroy when empty.

This granular product is for dry application on non-cropland where bare ground is desired. It is an effective general herbicide for the control of many annual and perennial grasses, weeds and brush. Effects are slow to appear and may not show until the chemical has been carried into the root zone by moisture. The degree of control obtained and length of control will vary with the amount of chemical applied, soil type, rainfall and other conditions. This product is non-corrosive to metals and non volatile.

IMPORTANT

Do not apply or drain or flush equipment on or near desirable trees or other plants, or areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of dry powder or spray to desirable plants. DO NOT contaminate domestic or irrigation waters. Keep from contact with fertilizers, insecticides, fungicides, and seeds. Thoroughly clean application equipment after use. DO NOT use equipment with which this product has been applied for applying insecticides, fungicides, and fertilizers as plant injury may result.

DIRECTIONS

Apply this product just before or during the period of active growth of weeds to be controlled. Moisture is necessary after treatment to carry the chemical into the root zone of weeds. Where dense growth is present results will be improved if vegetation is removed before treatment. Do not apply on frozen ground.

Any of the usual spreaders for granular materials should be acceptable for use in applying this product. Be sure to check calibration.

GENERAL WEED CONTROL

For an extended period of time most weeds may be controlled on non-cropland areas such as railroad and utility rights of way, lumber yards, pole and other storage yards, industrial sites, and tank farm.

To control perennial grasses such as quack grass, bluegrass, smooth brome grass apply $3\frac{1}{2}$ pound to $13\frac{1}{2}$ pounds per 100 square feet. For hard to kill species such as Johnson grass, Bermuda grass and nut grass apply $11\frac{1}{2}$ pounds to $21\frac{1}{2}$ pounds per 100 square feet.

To control perennial broad leafed weeds such as dock, wild carrot, sheep sorrel, plantain and dandelion, apply $11\frac{1}{2}$ to $13\frac{1}{2}$ pounds per 100 square feet. For hard to kill species such as brome grass, spurge, dogbane, goldenrod, daisies and asters, apply $11\frac{1}{2}$ pounds to $21\frac{1}{2}$ pounds per 100 square feet.

To control annual weeds and grasses such as foxtail, crabgrass, chickweed, wild oats, brome grass, ryegrass, pigweed, ragweed, purslane, wild mustard, cocklebur and lambsquarter, apply $3\frac{1}{2}$ to 10 pound per 100 square feet.

Use the higher application rates on those soils with high organic matter or high carbon levels.

BRUSH CONTROL

GRANULAR WEED AND BRUSH KILLER

ACTIVE INGREDIENTS:

Bromacil
(5-Bromo-3-sec-butyl-6-methyluracil) 2.00%

INERT INGREDIENTS 98.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN. May cause irritation of eyes, nose, throat and skin.
Avoid breathing dust. Avoid contact with skin, eyes, and clothing.

Do not reuse container. Destroy when empty.

This granular product is for dry application on non-cropland where bare ground is desired. It is an effective general herbicide for the control of many annual and perennial grasses, weeds and brush. Effects are slow to appear and may not show until the chemical has been carried into the root zone by moisture. The degree of control obtained and length of control will vary with the amount of chemical applied, soil type, rainfall and other conditions. This product is non-corrosive to metals and non-volatile.

IMPORTANT

Do not apply or drain or flush equipment on or near desirable trees or other plants, or areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of dry powder or spray to desirable plants. DO NOT contaminate domestic or irrigation waters. Keep from contact with fertilizers, insecticides, fungicides, and seeds. Thoroughly clean application equipment after use. DO NOT use equipment with which this product has been applied for applying insecticides, fungicides, and fertilizers as plant injury may result.

DIRECTIONS

Apply this product just before or during the period of active growth of weeds to be controlled. Moisture is necessary after treatment to carry the chemical into the root zone of weeds. Where dense growth is present results will be improved if vegetation is removed before treatment. Do not apply on frozen ground.

Any of the usual spreaders for granular materials should be acceptable for use in applying this product. Be sure to check calibration.

GENERAL WEED CONTROL

For an extended period of time most weeds may be controlled on non-cropland areas such as railroad and utility rights of way, lumber yards, pole and other storage yards, industrial sites, and tank farms:

To control perennial grasses such as quack grass, bluegrass, smooth brome grass apply $3\frac{1}{2}$ pound to $13\frac{1}{2}$ pounds per 100 square feet. For hard to kill species such as Johnson grass, Bermuda grass and nut grass apply $11\frac{1}{2}$ pounds to $21\frac{1}{2}$ pounds per 100 square feet.

To control perennial broad leafed weeds such as dock, wild carrot, sheep sorrell, plantain and dandelion, apply $11\frac{1}{2}$ to $13\frac{1}{2}$ pounds per 100 square feet. For hard to kill species such as bouncing bet, spurge, dogbane, goldenrod, daisies and asters, apply $11\frac{1}{2}$ pounds to $21\frac{1}{2}$ pounds per 100 square feet.

To control annual weeds and grasses such as foxtail, crabgrass, cheat grass, wild oats, brome grass, rye grass, pigweed, ragweed, purslane, wild mustard, cocklebur and lambsquarter, apply $\frac{1}{2}$ to 1 pound per 100 square feet.

Use the higher application rates on those soils with high organic matter or high carbon levels.

BRUSH CONTROL

Use as a spot treatment for control of willow, cottonwood, soft maple and certain other undesirable woody species in non-cropland areas. Effects are slow to appear and will not appear until the chemical has been activated by moisture. Apply during the spring and summer months.

Use as a spot treatment for control of willow, cottonwood, soft maple and certain other undesirable woody species in non-cropland areas. Effects are slow to appear and will not appear until the chemical has been activated by moisture. Apply during the spring and summer months.

DO NOT apply to brush standing in water.

NOTICE OF WARRANTY

Crown Chemicals, Inc. makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

MANUFACTURED BY

EPA Reg. No. 133-100

CROWN CHEMICALS

ROCKFORD, ILLINOIS 61101