

[7273-114]

CROWN

#### FARM USES

**MILKHOUSES AND DAIRY ROOMS** - Spray windows, doors, ceilings as needed but do not apply directly on foods and utensils.

**BARNs** - Spray resting places such as windows, doorways, ceilings where flies and mosquitoes congregate.

Spray ceilings and walls of animal pens to reduce fly breeding.

#### FOOD PROCESSING AREAS OF FOOD PLANTS, RESTAURANTS & HOME USES

For use in Food Processing Areas of Food Plants and Restaurants and Homes, **OBSERVE THE FOLLOWING CAUTIONS:**

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when plant or restaurant is not in operation.

**FOR CONTROL OF FLYING INSECTS** aim spray upward and distribute throughout space being treated. For maximum effectiveness keep doors and windows closed for 15 minutes after treatment.

**CRAWLING INSECTS** - Treat infested areas thoroughly and repeat as necessary, hitting as many insects as possible with the spray.

**FLIES, MOSQUITOES, GNATS, WASPS, HORNETS** as indicated for flying insects.

**ROACHES, ANTS, SILVERFISH, WATERBUGS** treat areas where the insects are commonly found, and direct spray behind baseboards, moldings, sinks and pipes, into cracks and crevices. Direct into locations where pests congregate and treat places where they enter.

**FLEAS, TICKS** - Apply to infested areas, thoroughly spraying all floors and floor coverings, and include treatment of sleeping quarters of pets. Repeat as necessary.

**CLOVER MITES, CRICKETS, SPIDERS, SCORPIONS** - Thoroughly spray floors of storage areas, closets, corners around pipes, and all areas where these pests are seen to crawl or congregate, spraying the insects directly whenever possible.

## FOOD PLANT & DAIRY ROOM INSECT BOMB



**SKIPPERS** - To aid in the control of exposed stages of skippers and the stored grain pests; granary weevil, rice weevil, confused flour beetle, rust red flour beetle, cadelle, saw-toothed grain beetle, spider beetle, angoumois grain moth, Mediterranean flour moth, meal worms, cigarette beetle, drug store beetle, and grain mite - Spray the outside of packaged cereals or food, spraying shelves, and other places where the pests are seen to alight, crawl and congregate, except on exposed food. Avoid sources of reinfestation by destroying wormy flour, grain and cereals before applications.

Do not remain in treated area. Ventilate the area after treatment is completed.

For Flies and Mosquitoes which enter through open doors, spraying a few seconds at a time at frequent intervals is effective.

#### WARNING

Harmful if swallowed. Avoid inhalation and skin contact. Remove pets, birds and cover fish aquariums before spraying. Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° Fahrenheit may cause bursting. Never throw container into fire or incinerator. Discard in a safe place.

#### NOTICE OF WARRANTY

Crown Chemicals makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

CROWN CHEMICALS  
Rockford, Illinois

E.P.A. Reg. No. 7273-114



# FOOD PLANT & DAIRY ROOM INSECT BOMB



## CONTAINS PYRETHRUM

For Control of

Roaches, Silverfish, Crickets,  
Spiders, Scorpions, Ants, Flies,  
Mosquitoes, Ticks, Gnats, Wasps,  
Hornets, Clover Mites, Fleas,  
Skippers, as well as stored grain  
pests:

### ACTIVE INGREDIENTS: By WEIGHT

Pyrethrins . . . . .	0.25%
Technical Piperonyl Butoxide . . .	1.25%*
Petroleum Distillates . . . . .	33.50%

INERT INGREDIENTS: . . . . . 65.00%

TOTAL . . . . . 100.00%

\*Equivalent to 1.0% (butylcarbityl) (6-propyl-  
piperonyl) ether and .25% other related com-  
pounds.

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
See Back Panel For Additional Precautions

NET CONTENTS 16 OZ.