

SECRET

100-443887-100



APPROVED
 [Signature]
 [Stamp]

PRECAUTIONS IN USING

DO NOT USE FOR STORES OF FOOD OR FEED. Do not use in areas where food or feed is stored. Do not use in areas where food or feed is stored. Do not use in areas where food or feed is stored.

DO NOT USE IN AREAS WHERE FOOD OR FEED IS STORED. Do not use in areas where food or feed is stored. Do not use in areas where food or feed is stored.

DANGER: Do not get on face, eyes, nose, mouth, or throat. If swallowed, do not induce vomiting. If swallowed, do not induce vomiting. If swallowed, do not induce vomiting.

This product is toxic to fish and wild life. Do not feed to domestic animals. Do not feed to domestic animals. Do not feed to domestic animals.

ANTIDOTE: If swallowed, give 1 tablespoon of 1% sodium bicarbonate solution with water and repeat every 15 minutes.

ANTIDOTE EXTERNAL: Remove contaminated clothing and wash with plenty of water. Flush with plenty of water. Do not use oil or grease.

CALL A PHYSICIAN IMMEDIATELY. Antidote is 1% sodium bicarbonate solution. Antidote is 1% sodium bicarbonate solution. Antidote is 1% sodium bicarbonate solution.

DIRECTIONS FOR USE

1. For control of house flies, apply 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

FACE FLY SOLUTION: Mix 1/2 ounce of Face Fly Solution with 1 gallon of water.

SOLUTION: Mix 1/2 ounce of Face Fly Solution with 1 gallon of water.

FOGGING SOLUTION: Mix 1/2 ounce of Face Fly Solution with 1 gallon of water.

DO NOT MIX UP MORE DILUTED SPRAY THAN IS NEEDED FOR THE APPLICATION.

FACE FLIES: Face Fly Solution with sugar. Apply 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

LACTATING DAIRY & BEEF CATTLE, HORSES: 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

HOGS, SHEEP AND GOATS: 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

FOGGING AND RESIDUAL ANIMAL BUILDINGS, DAIRY BARN, MILK ROOMS, STABLES, HOG HOUSES, SHELTERS AND POULTRY HOUSES: 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

DO NOT CONTAMINATE MILK, WATER, FEED OR UTENSILS.

FLY BREEDING AREAS, MANURE PILES, GARBAGE DUMPS: 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

FEED LOTS, BEEF & HOG STOCKYARDS FOR CONTROL OF FLIES, MOSQUITOES, GNATS: 1/2 ounce of Face Fly Solution per 1000 cubic feet of space.

DO NOT FEED TO DOMESTIC ANIMALS.

NOTICE OF WARRANTY: This product is warranted to be free from defects in material and workmanship.