




DIPHACINONE CONCENTRATE RODENTICIDE

Anti-Coagulant For Rodent Control

CAUTION

KEEP OUT OF THE REACH OF CHILDREN
See Below For Additional Precautions

ACTIVE INGREDIENT:

Diphacinone (2-diphenylacetyl-1,3-indandione) 0.1 %

INERT INGREDIENT 99.9 %

Total 100.0 %

GENERAL DIRECTIONS

- a. Do not use on or near water, or on or near the water supply, or on or near the water supply, or on or near the water supply.
- b. Place bait in areas where rodents are known to travel, and where they are not likely to be disturbed by other animals.
- c. Bait placement should be made at night, and should be made in areas where rodents are known to travel, and where they are not likely to be disturbed by other animals.
- d. Place bait in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.

ACTIVE INGREDIENT:

Diphacinone (2-diphenylacetyl-1,3-indandione) 0.1%

INERT INGREDIENT 99.9%

Total 100.0%

GENERAL DIRECTIONS

- a) Thoroughly mix equal parts by weight of diphacinone with nine parts of dry cereal bait.
- b) Place bait in locations where rats and mice feed, water and/or travel.
- c) Bait placements along walls, in corners, and in concealed locations where rats and mice can feed undisturbed are desirable.
- d) Place baits in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes.

DIRECTIONS FOR RAT CONTROL

- a) Place 4 to 16 ounces of bait at each bait location.
- b) Maintain an uninterrupted supply of bait in the stations for at least 10 days.

DIRECTION FOR MOUSE CONTROL

- a) Place bait (specimens of 4 to 12 ounces) at bait at 8 to 12 foot intervals.
- b) Maintain an uninterrupted supply of bait for at least 15 days.

NOTE: If bait becomes stale or becomes hard, replace with fresh material.

CAUTION

Keep away from human consumption areas, wildlife and pets. If swallowed by human or animal, it may irritate the stomach and may reduce the clotting ability of the blood, and cause bleeding. In such cases, the intravenous and/or oral administration of Vitamin K is indicated, the physician should be immediately notified in the case of human cases caused by ingestion of this product.

Do not use contents of this container for any other purpose.

© 1977 Crown Chemicals

CROWN CHEMICALS
Rockford, Illinois 61101