



#### DIRECTIONS FOR USE

Hold container at a distance from the area to be treated that will permit even distribution of the dust. Keep lid closed when container is not in use.

#### RECOMMENDED APPLICATIONS

ANIMAL	PARASITE	REMARKS
Cattle (Dairy and Beef)	Horn Flies Lice	<b>DIRECT APPLICATION:</b> Apply not more than 2 oz.* /animal. Dust evenly over the head, neck, shoulders, back and tailhead. Repeat as necessary. No interval is required between treatment and use of meat or milk for food.
Swine	Lice	<b>DIRECT APPLICATION:</b> Apply not more than 1 oz.* /animal as a uniform coat to the shoulders and back. Repeat as necessary but not more often than once every 10 days. No interval is required between treatment and use of meat as food.  <b>BEDDING TREATMENT:</b> Apply 2 oz.* uniformly over each 30 square feet of fresh, dry bedding. Repeat as necessary, but not more than once every 10 days.  <b>DIRECT APPLICATION AND BEDDING TREATMENT:</b> In severe infestations, both individual animals and the bedding may be treated as directed above.


\* Note: 1 oz. of this material equals approximately three level tablespoonsful.

#### RESTRICTIONS

For external, insecticidal use on above specified animals only. Avoid contamination of feed, troughs, water and water utensils. Provide thorough ventilation while dusting.

# 'D-LOUSE'<sup>TM</sup>

## ANIMAL INSECTICIDE



#### CAUTION

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

#### CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

TO PHYSICIAN — Atropine sulfate is antidotal. 2-PAM is also antidotal, and may be administered in conjunction with atropine.

Do not use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

#### NOTICE OF WARRANTY

Crown Chemicals, Inc. makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

Manufactured for  
**CROWN CHEMICALS**  
ROCKFORD, ILLINOIS 61101

EPA Reg. No. 7273-127

7273-127

11-8-1972

1/2

ACCEPTED

NOV 8 1972

ORDER TO TEST AN INSECTICIDE  
FOR USE IN AND A PESTICIDE ACT  
FOR AN AGRICULTURAL PESTICIDE  
REGISTERED UNDER NO. 1213-121 SUBJECT  
TO ATTACHED COMMENTS.