

"PEST RID"

PELLETED BAIT KILLS RATS AND MICE

Each particle of the warfarin in this product has been specially coated with a material that reduces detection of the warfarin by the rodents at the time the bait is consumed. The coating may help assure greater acceptance of the bait in cases where rodent awareness of uncoated warfarin is responsible for low bait acceptability.

COTNE INGREDIENT

Lastain 3- Spins & Morgan Co.

Fig. 1. Continues and the second of the seco

Testeri

CAUTION KEEP OUT OF REACH OF CHILDREN

See Additional Precautions Below.

INSTRUCTIONS

RATS Open package and place four ounces of "Pest Rid" in dry ontainer next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of built for a period of not less than fifteen days and continue builting until all signs of feeding have stopped. Replace state, damp or holdy baits with fresh bait.

MICE Open package and place tablespoonful amounts of "Perr Rid" bart in shallow containers where mice or signs of nice have been seen. Bart stations at intervals of eight to twelve feet are recommended. Assure an ununterrupted copply of bart for a new of eight iess than fifteen day.

The Complete of the constants want be constants and a transfer of a section of the constant of

awareness of uncoated warrarin is responsible for low bait acceptability.

ALTER INGREDIENT

We with the an exception of express they got

Frank Moory on many the contract of the contra The state of the state of the state of

7 11 11 111

CAUTION KEEP OUT OF REACH OF CHILDREN

See Additional Precautions Below.

INSTRUCTIONS

RATS: Open package and place four ounces of "Pest Rid" in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than fifteen days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

Open package and place tablespoonful amounts of "Pest Rid" bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of eight to twelve feet are recommended. Assure an uninterrupted supply of pait for a period of not less than fifteen days.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label

DO NOT Reuse Container Destroy When Empty

If swallowed by humans, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of Bishydroxycoumarin In case of ingestion call a physician

NET WT

Manufactured by

CROWN CHEMICALS / , Rockford Illinois 61101