

DIRECTIONS FOR USE HOUSEFLIES

Residual Wall Sprays For the control of house flies, including resistant strains in dairy farms, hog pens, and barns, poultry houses, and other farm buildings, apply a 1% residual spray to the ceilings, walls, stanchions, etc. Prepare the spray by mixing 1 pint Flyban 2-E in 3 gallons of water. Thoroughly wet all fly resting areas to the point of run off. One gallon of spray will cover 500 to 1000 square feet of surface. Repeat applications should be made when necessary.

Remove all animals from buildings when apply residual wall sprays.

Flyban 2-E is acceptable for use as a residual-type insecticide in and around federally inspected meatpacking plants except where meat food products are processed or handled.

Flyban 2-E controls flies up to 8 weeks or longer.

For Localized Housefly Control Apply a spray containing 1/2 pint of Flyban 2-E in 5 quarts of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, etc.

Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

Maggot Sprays For the control of housefly maggots, mix 1/2 pint of Flyban 2-E in 5 quarts of water, and apply as a coarse spray or with a sprinkling can to fly-breeding areas, such as poultry droppings in caged layer houses, garbage dumps and manure piles. Repeat application as additional manure or garbage is added.

General Outside Use For the control of house flies around homes and recreation areas, garbage cans, animal quarters, food processing plants, warehouses, loading docks and refuse areas, thoroughly spray exposed surfaces, such as walls, fences, garbage and refuse containers with 1 pint of Flyban 2-E in 3 gallons of water. Repeat applications should be made when necessary.

DO NOT Contaminate Feed and Foodstuffs, Drinking Fountains, Litter and Feed Troughs.

DO NOT use in Milk Processing Rooms, including Milk Houses and Milk Storage Rooms.

DO NOT use in Homes.

DO NOT use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas where food is exposed.

CROWN

FLYBAN 2-E

Systemic Insecticide

ACCEPTED

7273-140

OCT 24 1974

USE IN ACCORDANCE WITH LABEL DIRECTIONS FOR THE CONTROL OF HOUSEFLIES AND MAGGOTS IN AND AROUND MEATPACKING PLANTS AND OTHER FACILITIES.

CONTROLS HOUSEFLIES

For control of houseflies and maggots in and around meatpacking plants and other facilities.

ACTIVE INGREDIENTS

Chlorpyrifos 20% (1.0% active)
Diazinon 10% (1.0% active)
Xylenol 1.0%

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Precautions and First Aid

7273-140

23-1L-1

NET CONTENTS

GALLONS

VAPOR
SWALLOWED
CAUTION
DO NOT
DO NOT
Use
Waste
DO NOT
DO NOT
open
In case
plem
medic
NOTE
chole
dota
DO NOT
TUNE
Rinse
When
taint
card
This
life
may
pond
occu
favo
nate
posi
spec
CRC
char
wise
duct
ment